

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 May 2003 (08.05.2003)

PCT

(10) International Publication Number
WO 03/037432 A1

(51) International Patent Classification⁷: **A61P 3/06**, 3/08,
3/10, 9/10, 9/12, **A61K 31/519**, 45/06

(74) Agents: **LUMB, J., Trevor** et al.; Pfizer Inc., 201 Tabor
Road, Morris Plains, NJ 07950 (US).

(21) International Application Number: PCT/IB02/03754

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(22) International Filing Date:
12 September 2002 (12.09.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/336,981 2 November 2001 (02.11.2001) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **PFIZER
PRODUCTS INC.** [US/US]; Eastern Point Road, Groton,
CT 06340 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FRYBURG, David,**
Albert [US/US]; Pfizer Global Research and Development,
Eastern Point Road, Groton, CT 06340 (US). **GIBBS, Earl,**
Michael [US/US]; Pfizer Global Research and Develop-
ment, Eastern Point Road, Groton, CT 06340 (US).

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 03/037432 A1

(54) Title: TREATMENT OF INSULIN RESISTANCE SYNDROME AND TYPE 2 DIABETES WITH PDE9 INHIBITORS

(57) Abstract: This invention is directed to a method of treating insulin resistance syndrome (IRS), hypertension and/or type 2 diabetes in a mammal comprising administering to said mammal a cGMP PDE9 inhibitor or a pharmaceutical composition thereof. This invention is also directed to such methods wherein said cGMP PDE9 inhibitor is used in combination with other agents to treat IRS, hypertension and/or type 2 diabetes.

BEST AVAILABLE COPY

TREATMENT OF INSULIN RESISTANCE SYNDROME
AND TYPE 2 DIABETES WITH PDE9 INHIBITORS

FIELD OF THE INVENTION

5 This invention relates to the use of cGMP PDE9 inhibitors for the treatment of type
2 diabetes, hyperglycemia, dyslipidemia, impaired glucose tolerance, type 1 diabetes
and/or insulin resistance syndrome (IRS). This invention also relates to combinations
comprising cGMP PDE9 inhibitors and other agents, said combinations being useful in
treating type 2 diabetes, hyperglycemia, dyslipidemia, impaired glucose tolerance, type 1
10 diabetes and/or insulin resistance syndrome.

BACKGROUND OF THE INVENTION

IRS, as defined herein, means the concomitant existence in a subject of two or
more of: hyperinsulinemia, dyslipidemia, hypertension, type 2 diabetes or impaired
glucose tolerance, hyperuricemia or gout, a pro-coagulant state, atherosclerosis and/or
15 truncal obesity. At the center of IRS, also known as "Syndrome X" and "Metabolic
Syndrome" in the biomedical literature, is the common feature of tissue resistance to the
action of insulin. This impaired biological response to insulin is manifested in the
metabolic and vascular effects of insulin. Although there are monogenic syndromes of
insulin resistance (IR), in which a definite gene has been identified as the cause of insulin
20 resistance (such as leprechaunism), these are relatively rare. By contrast, the more
common presentation of the IRS is associated with obesity (particularly abdominal) and
appears to be polygenic.

The adaptive response to IR in individuals having IRS produces compensatory
hyperinsulinaemia. As subjects with IRS become progressively insulin resistant, they
25 manifest varying degrees of change in clinical parameters, including blood pressure,
and/or increased levels of serum glucose, and/or cholesterol and/or triglycerides, and/or
uric acid, and/or factors that increase coagulation. Once these clinical parameters have
changed enough, the patient with IRS may differentially manifest well-recognized clinical
conditions or diagnoses.

30 2. These conditions include: type 2 diabetes, hypertension (high blood
pressure), hyperlipidemia or dyslipidemia, particularly (but not limited to)
hypertriglyceridemia, hyperuricemia or gout, and hypercoagulability (defined as an
abnormal, increased tendency for clots to form, particularly inside blood vessels). These
clinical conditions are well-recognized risk factors for cardiovascular (coronary artery and
35 cerebrovascular) disease.

While it is difficult to estimate the prevalence of IRS in the general populace due to both the diversity of the collective risk factors associated with the syndrome and the likelihood that many individuals affected by IRS go undetected because they may exhibit no exterior symptoms and have no prior history of coronary heart disease, it is postulated that at a minimum the patient population at risk for the development of IRS includes individuals with obesity, particularly truncal (abdominal) obesity. Obesity is an extremely common problem in the industrialized world and is associated with the clinical conditions mentioned above. Thus, it is very likely that the prevalence of IRS is very high. Considering this potential patient group alone forms an immense population potentially at risk for the development of complications of IRS. For example in the United States in 1994, 23% of the population aged between 20 and 74 had hypertension, which accounted for 5 deaths per 100,000 population (1997). There will be an estimated 154,392,000 patients with diabetes world-wide in the year 2000. Of these, 15,000,000 will be in the US and 934,000 in the UK. The burden of disease for ischaemic heart disease for both sexes in the WHO region estimated for 1998 was 51,948,000 with a mortality of 7,375,000, constituting 13.7% of total mortality and ranking the highest in the mortality score. The burden of diabetes in both sexes in the WHO region estimated for 1998 was 11,668,000. Thus there exists a large medical need for an effective and safe oral therapy for the treatment of IRS and prevention of the development of IRS and its clinical consequences.

Resistance to the effects of insulin is also observed in the diminished biological response of the endothelium to the vascular effects of insulin. That is, insulin promotes relaxation of blood vessels at least in part through the action of nitric oxide (NO). Nitric oxide generated in the endothelium then stimulates cGMP production in blood vessels and causes them to relax or dilate. This opening of the blood vessel allows more blood to flow, which is particularly important when more blood flow is needed to critical organs, like the heart. It has been demonstrated that there is a decreased release of NO from the endothelium of patients with IR. This decreased release of NO is not only from insulin, but also from other important vasodilators like acetylcholine. This so-called "endothelial dysfunction" contributes to the risk factors for cardiovascular disease which are associated with IRS. The vascular effect of insulin contributes to the effect of insulin to regulate metabolism, particularly, but not necessarily limited to, glucose metabolism.

NO also has direct effects on glucose uptake by skeletal muscle. That is, treatment with a NO-donor substance, such as nitroprusside, or with an analogue of cGMP *in vitro* increases glucose uptake (transport by GLUT4 glucose transporters). This vasodilation-independent pathway is described in G. J. Etgen, D. A. Fryburg and E. M.

Gibbs in *Diabetes*, 46, 1997 pp. 1915-1919, which is incorporated herein by reference. Taken together, NO and cGMP have direct target tissue (skeletal muscle) and vascular actions that influence, mediate, or mimic the action of insulin.

Further effects of impaired NO release by the endothelium include: increases in
5 vascular smooth muscle cell (VSMC) growth, proliferation and migration which are key steps in atherosclerotic plaque formation that can lead to stroke; an increase in platelet aggregation and adhesiveness; an increase of lipid peroxidation and an effect on the inhibition of cell adhesion molecule expression including vascular cell adhesion molecule (VCAM-1), intracellular adhesion molecule (ICAM), E-selectin. Impaired endothelial NO
10 release also impacts on the activity of inflammatory cytokines such as tumour necrosis factor- α (TNF- α), and the production of monocyte chemoattractant factor through decreased activity of the transcriptional activator nuclear factor kappa B. These effects on the platelet are also cGMP driven.

Finally, there are examples in which the treatment of factors contributing to IRS
15 (e.g., obesity) or the treatment of IRS itself improves many of these clinical conditions which at first glance appear to be unrelated. For example, dieting alone or pharmacotherapeutic agents that induce weight loss will decrease blood pressure, blood glucose and triglycerides. Agents that are designed to improve insulin sensitivity can also favorably alter blood pressure, lipids, and blood glucose.

20 Successful diagnosis and treatment of patients with IRS with a PDE9 inhibitor will lead to clinically relevant improvements in blood pressure, and/or serum glucose and/or insulin and/or lipids and/or uric acid, and/or procoagulant factors. This treatment can occur alone or in combination with other therapeutics that improve IRS. Improvement in these clinical conditions should reduce the risk of the development of cardiovascular
25 disease in these patients as well as other complications of these individual disorders (including, but not limited to diabetic neuropathy, nephropathy, and retinopathy).

While IRS has many manifestations, an important underlying mechanistic basis for the condition resides in a resistance to both the vascular and metabolic effects of insulin. It is also understood that the underlying pathology of vascular resistance in insulin
30 resistance syndrome, is a diminished amount of NO produced by the endothelial cells in response to insulin. There is impaired signaling of insulin for glucose uptake in insulin resistant individuals.

Amplification of the cGMP signal, using cGMP PDE9 inhibitors in patients with IRS enhances the insulin glucose uptake signal and improves insulin action at key tissues.
35 Enhancing insulin sensitivity improves clinical parameters of IRS results, *inter alia*, in:

1. Blood glucose control: In patients with type 2 diabetes or impaired glucose tolerance, an improvement in insulin sensitivity results in a decrease in plasma glucose concentrations (either fasting or after an oral glucose tolerance test or a meal). In a related manner, as regulated by the patient's pathophysiology, there will be an improvement in serum insulin concentrations in either the fasting state or after a glucose load or meal. These improvements in blood glucose control, should the subjects have type 2 diabetes, manifest as improvements in measures of long-term blood glucose control, such as, but not limited to, hemoglobin A1c (glycosylated hemoglobin) or fructosamine.
2. Blood pressure: It is believed that improvement in insulin sensitivity yields improvements in both systolic and diastolic blood pressure.
3. Lipids: Improvement in insulin resistance yields improvements in serum lipids, including, but not limited to, serum cholesterol and triglycerides.
4. Uric Acid: Improvement in insulin resistance yields improvements in serum uric acid.
5. Coagulation Factors: It is believed that improvement in insulin resistance restores normal factors that worsen the procoagulant state.

cGMP PDE 9 inhibitors prevent the effect of the phosphodiesterase 9 enzyme that converts cGMP to inactive GMP thus increasing the amount of accumulated cGMP. This accumulation amplifies the vasodilatory, metabolic, and anti-atherogenic effects of the available nitric oxide and insulin. This amplification action mitigates the adverse effects associated with IRS and improve one or more of the associated conditions.

Diabetes mellitus is characterized by metabolic defects in production and utilization of carbohydrates, resulting in elevated blood glucose or hyperglycemia due to the failure to maintain appropriate blood sugar levels. Research in the treatment of diabetes has centered on attempts to normalize fasting and postprandial blood glucose levels. Current treatments include administration of exogenous insulin, oral administration of drugs and dietary therapies and exercise regimens.

Two major forms of diabetes mellitus are recognized. Type 1 diabetes, or insulin-dependent diabetes, is the result of an absolute deficiency of insulin, the hormone which regulates carbohydrate utilization. Type 2 diabetes, or non-insulin dependent diabetes, often occurs with normal, or even elevated levels of insulin and appears to be the result of the inability of tissues to respond appropriately to insulin. Complications of type 2 diabetes include retinopathy, nephropathy, neuropathy, and coronary heart disease, and are believed to be triggered by excessive protein glycation, which in turn results from excessive levels of circulating glucose. Reduction in hyperglycemia by treatment with a

PDE9 inhibitor will lower the level of protein glycation and result in a diminution in these diabetic complications.

Polycystic ovary syndrome (PCOS) also known as Stein-Leventhal syndrome or functional ovarian hyperandrogenism, is a complex endocrine disorder associated with a long-term lack of ovulation (anovulation) and an excess of androgens (male sex hormones, e.g., testosterone) circulating in the blood. The disorder is characterized by the formation of cysts in the ovaries, a process related to the failure of the ovary to release an egg (ovum). In the majority of cases, the ovaries become enlarged. PCOS afflicts up to 22% of women during their childbearing years, although only 10% of these women develop symptoms. It is one of the most frequent causes of infertility in women.

SUMMARY OF THE INVENTION

This invention is directed to a method of treating IRS in a mammal comprising administering to said mammal a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said PDE9 inhibitor, prodrug or solvate. In a preferred embodiment of this invention, said method comprises administering a pharmaceutical composition comprising a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said PDE9 inhibitor, prodrug or solvate. Preferably, said pharmaceutical composition additionally comprises a pharmaceutically acceptable vehicle, diluent or carrier.

This invention is also directed to a method of treating type 2 diabetes in a mammal comprising administering to said mammal a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said PDE9 inhibitor, prodrug or solvate. In a preferred embodiment of this invention, said method comprises administering a pharmaceutical composition comprising a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said PDE9 inhibitor, prodrug or solvate. Preferably, said pharmaceutical composition additionally comprises a pharmaceutically acceptable vehicle, diluent or carrier.

This invention is also directed to a method of treating type 1 diabetes in a mammal comprising administering to said mammal a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said cGMP PDE9 inhibitor, prodrug or solvate. In a preferred embodiment of this invention, said method comprises administering a pharmaceutical composition comprising a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said cGMP PDE9 inhibitor, prodrug or solvate. Preferably, said pharmaceutical composition additionally comprises a pharmaceutically acceptable vehicle, diluent or carrier.

This invention is also directed to a method of treating impaired glucose tolerance

in a mammal comprising administering to said mammal a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said cGMP PDE9 inhibitor, prodrug or solvate. In a preferred embodiment of this invention, said method comprises administering a pharmaceutical composition comprising a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said cGMP PDE9 inhibitor, prodrug or solvate. Preferably, said pharmaceutical composition additionally comprises a pharmaceutically acceptable vehicle, diluent or carrier.

This invention is also directed to a method of treating dyslipidemia such as, but not limited to, hypertriglyceridemia and high LDL cholesterol, in a mammal comprising administering to said mammal a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said cGMP PDE9 inhibitor, prodrug or solvate. Dyslipidemia, where used herein, means an alteration of the lipid profile in blood.. In a preferred embodiment of this invention, said method comprises administering a pharmaceutical composition comprising a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said cGMP PDE9 inhibitor, prodrug or solvate. Preferably, said pharmaceutical composition additionally comprises a pharmaceutically acceptable vehicle, diluent or carrier.

This invention is also directed to a method of treating polycystic ovary syndrome in a mammal comprising administering to said mammal a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said cGMP PDE9 inhibitor, prodrug or solvate. In a preferred embodiment of this invention, said method comprises administering a pharmaceutical composition comprising a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said cGMP PDE9 inhibitor, prodrug or solvate. Preferably, said pharmaceutical composition additionally comprises a pharmaceutically acceptable vehicle, diluent or carrier.

In a further embodiment, this invention is directed to a first combination comprising two active ingredients selected from a cGMP PDE9 inhibitor, a prodrug, solvate or salt thereof and one or more, independently selected, protein kinase inhibitor, prodrug, solvate or salt thereof; an AMP-activated protein kinase activator, prodrug, solvate or salt thereof; a weight loss agent, prodrug, solvate or salt thereof; insulin; a PPAR- γ agonist, prodrug, solvate or salt thereof; a PPAR- γ antagonist, prodrug, solvate or salt thereof, a PPAR- α agonist, prodrug, solvate or salt thereof; a dual PPAR- γ /PPAR- α agonist, prodrug, solvate or salt thereof; a sorbitol dehydrogenase inhibitor, prodrug, solvate or salt thereof; a glycogen phosphorylase inhibitor, prodrug, solvate or salt thereof; a biguanide such as metformin, prodrug, solvate or salt thereof; an HMG-CoA reductase inhibitor, prodrug, solvate or salt thereof; an aldose reductase inhibitor, prodrug, solvate

or salt thereof; a PDE5 inhibitor, prodrug, solvate or salt thereof; a PDE11 inhibitor, prodrug, solvate or salt thereof; or a CETP inhibitor, prodrug, solvate or salt thereof. An especially preferred combination is a combination of a cGMP PDE9 inhibitor, a prodrug, solvate or salt thereof and a PDE5 inhibitor, a prodrug, solvate or salt thereof. In a further
5 embodiment, the invention is directed to a pharmaceutical composition comprising said first combination and a pharmaceutically acceptable vehicle, carrier or diluent. In a further embodiment, this invention is directed to methods of treating insulin resistance in a mammal comprising administering to said mammal said first combination or a pharmaceutical composition comprising said first combination. In a still further
10 embodiment, this invention is directed to a method of treating type 2 diabetes in a mammal comprising administering to said mammal said first combination or a pharmaceutical composition comprising said first combination.

In a still further embodiment, this invention is directed to a second combination comprising three active ingredients selected from a cGMP PDE9 inhibitor, a prodrug, solvate or salt thereof; a cGMP PDE5 inhibitor, a prodrug, solvate or salt thereof; and a
15 cGMP PDE11 inhibitor, a prodrug, solvate or salt thereof. In a further embodiment, the invention is directed to a pharmaceutical composition comprising said second combination and a pharmaceutically acceptable vehicle, carrier or diluent. In a further embodiment, this invention is directed to methods of treating insulin resistance in a mammal comprising administering to said mammal said second combination or a pharmaceutical composition comprising said second combination. In a still further
20 embodiment, this invention is directed to a method of treating type 2 diabetes in a mammal comprising administering to said mammal said second combination or a pharmaceutical composition comprising said second combination.

25 This invention is also directed to a kit comprising:

- a) a first unit dosage form comprising a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said compound, prodrug or solvate and a pharmaceutically acceptable carrier, vehicle or diluent;
- b) a second unit dosage form comprising:
30 a protein kinase inhibitor;
an AMP-activated protein kinase;
a weight loss agent;
insulin;
a PPAR- γ agonist;
35 a PPAR- γ antagonist;
a PPAR- α agonist;

- a dual PPAR- γ /PPAR- α agonist;
a sorbitol dehydrogenase inhibitor;
a glycogen phosphorylase inhibitor;
a biguanide such as metformin
an HMG-CoA reductase inhibitor;
an aldose reductase inhibitor;
a PDE5 inhibitor;
a PDE11 inhibitor; or
a CETP inhibitor;
- 10 a prodrug or solvate of said protein kinase inhibitor, AMP-activated protein, weight loss agent, insulin, PPAR- γ agonist, PPAR- α agonist, PPAR- α antagonist, dual PPAR- γ /PPAR- α agonist, sorbitol dehydrogenase inhibitor, glycogen phosphorylase inhibitor, biguanide, vastatin, aldose reductase inhibitor, PDE5 inhibitor, PDE11 inhibitor or CETP inhibitor; or a pharmaceutically acceptable salt thereof or of said prodrug or solvate and a
- 15 pharmaceutically acceptable carrier, vehicle or diluent; and
- c) a container.

This invention is also directed to a kit comprising:

- a) a first unit dosage form comprising a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said compound,
- 20 prodrug or solvate and a pharmaceutically acceptable carrier, vehicle or diluent;
- b) a second unit dosage form comprising a cGMP PDE5 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said compound, prodrug or solvate and a pharmaceutically acceptable carrier, vehicle or diluent;
- c) a third unit dosage form comprising a cGMP PDE11 inhibitor, a
- 25 prodrug or solvate thereof or a pharmaceutically acceptable salt of said compound, prodrug or solvate and a pharmaceutically acceptable carrier, vehicle or diluent; and
- d) a container.

A further aspect of the invention provides for a method for treating IRS as defined above in a polygenic insulin resistant mammal comprising administering to the mammal

30 an effective amount of a cGMP PDE9 inhibitor or a pharmaceutical composition thereof. It is a further aspect of this invention to treat such polygenic insulin resistant mammals with a combination of a cGMP PDE9 inhibitor and a second compound as defined above, or with a pharmaceutical composition comprising such a combination and a pharmaceutically acceptable vehicle, carrier or diluent. In yet another aspect of this

35 invention, such polygenic insulin resistant mammals are treated with a kit as described above.


The suitability of any particular cGMP PDE9 inhibitor can be readily determined by evaluation of its potency and selectivity using literature methods followed by evaluation of its toxicity, absorption, metabolism, pharmacokinetics, etc in accordance with standard pharmaceutical practice.

- 5 Preferably, the cGMP PDE9 inhibitors have an IC_{50} at less than 100 nanomolar, more preferably, at less than 50 nanomolar, more preferably still at less than 15 nanomolar.

IC_{50} values for the cGMP PDE9 inhibitors may be determined using the PDE9 assay in the Test Methods Section hereinafter.

- 10 It is to be understood that the contents of the above published patent applications, and in particular the general formulae and exemplified compounds therein are incorporated herein in their entirety by reference thereto.

A preferred group of cGMP PDE9 inhibitors for use in the methods, compositions, combinations and kits of the instant invention include compounds of the formula (I)



15


or a pharmaceutically acceptable salt, solvate or prodrug thereof, wherein:

R^1 is H or (C_1-C_6) alkyl;

R^2 is (C_1-C_6) alkyl, straight chain or branched chain, (C_3-C_7) cycloalkyl or heteroaryl;

- 20 R^3 is (C_1-C_6) alkyl, straight chain or branched chain, optionally substituted by 1-2 groups each independently selected from Ar, (C_3-C_7) cycloalkyl, OAr, SAr, $NC(O)(C_1-C_6)$ alkyl, heteroaryl, xanthene, and naphthalene;

Ar is a group of formula



- 25 wherein R^4 , R^5 and R^6 are each independently selected from H, halo, phenoxy, phenyl, CF_3 , OCF_3 , $S(C_1-C_6)$ alkyl, (C_1-C_6) alkyl, $O(C_1-C_6)$ alkyl, said alkyl optionally substituted by a heteroaryl group or by a phenyl group, wherein said phenyl group is optionally substituted by 1-3 groups selected from halo, CF_3 , OCF_3 and (C_1-C_6) alkyl; or wherein R^4 and R^5 may combine to form a (C_2-C_3) alkyl link, wherein said link may optionally incorporate a
- 30 heteroatom selected from O, S and N; and

heteroaryl is aromatic 5-6 membered heterocycle containing 1-3 heteroatoms, each independently selected from O, S and N, said heterocycle optionally substituted by 1-3 substituents, each independently selected from (C₁-C₆)alkyl, halo and phenyl, said phenyl optionally substituted by 1-3 groups selected from halo and (C₁-C₆)alkyl;

- 5 with the proviso that when R¹ is -CH₃, R² cannot be -CH₂CH₂CH₃.

A particularly preferred group of compounds within the preferred group are those compounds wherein R¹ is H or CH₃. More preferably R¹ is H.

Another particularly preferred group of compounds within the preferred group are those compounds wherein R² is selected from (C₃-C₄)alkyl, cyclopentyl and pyridinyl.

- 10 More preferably R² is 3-pyridinyl.

Another particularly preferred group of compounds within the preferred group are those compounds wherein R³ is (C₁-C₃)alkyl, optionally substituted by 1-2 groups selected from: Ar, (C₃-C₇)cycloalkyl and heteroaryl. More preferably R³ is (C₁-C₃)alkyl, optionally substituted by Ar. Most preferably R³ is C₁ alkyl substituted by Ar, where R⁴, R⁵ and R⁶

15 are each H.

Another particularly preferred group of compounds within the preferred group are those compounds wherein R⁴, R⁵ and R⁶ are each independently selected from: H, halo, phenoxy, phenyl, CF₃, OCF₃, S(C₁-C₆)alkyl, (C₁-C₆)alkyl, O(C₁-C₆)alkyl, said alkyl optionally substituted by a heteroaryl group or by a phenyl group, wherein said phenyl

20 group is optionally substituted by 1-3 groups selected from halo, CF₃, OCF₃ and (C₁-C₆)alkyl; or wherein R⁴ and R⁵ may combine to form a C₂ alkyl link, said link incorporating an O atom. More preferably R⁴, R⁵ and R⁶ are each independently selected from H, halo, OCF₃, CF₃, OAr, and O(C₁-C₆)alkyl optionally substituted by phenyl, optionally substituted by H, halo, phenoxy, phenyl, CF₃, OCF₃, S(C₁-C₆)alkyl, (C₁-C₆)alkyl, O(C₁-C₆)alkyl, said

25 alkyl optionally substituted by a heteroaryl group or by a phenyl group, wherein said phenyl group is optionally substituted by H, halo, CF₃, OCF₃ and (C₁-C₆)alkyl. Yet more preferably R⁴, R⁵ and R⁶ are each independently selected from Cl, H, OCF₃, CF₃ and O(C₁-C₆)alkyl substituted by phenyl. Most preferably, R⁴, R⁵ and R⁶ are each independently selected from H, Cl and O(C₁-C₃)alkyl substituted by phenyl.

- 30 Another particularly preferred group of compounds within the preferred group are those compounds wherein heteroaryl is an aromatic 5-6 membered heterocycle containing at least 2 nitrogen atoms, said heterocycle optionally substituted by 1-3 substituents, each independently selected from (C₁-C₆)alkyl, halo and phenyl, said phenyl optionally substituted by 1-3 groups selected from halo and (C₁-C₆)alkyl. More preferably
- 35 heteroaryl is an aromatic 5 membered heterocycle containing at least 2 nitrogen atoms, said heterocycle optionally substituted by 1 substituent, each independently selected from

(C₁-C₆) alkyl, halo and phenyl, said phenyl optionally substituted by 1-3 groups selected from halo and (C₁-C₆) alkyl. Yet more preferably heteroaryl is an aromatic 5 membered heterocycle containing at least 2 nitrogen atoms, said heterocycle optionally substituted by phenyl optionally substituted by halo. Most preferably heteroaryl is an imidazole or an oxadiazole.

An especially preferred cGMP PDE9 inhibitor is 5-(3-chloro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one, a prodrug thereof or a pharmaceutically acceptable salt thereof or of said prodrug.

According to a further aspect the present invention additionally provides for the use of a PDE9 inhibitor or a pharmaceutical composition thereof for the treatment of the insulin resistance syndrome in a subject having type 2 diabetes mellitus or impaired glucose tolerance or having a family history of diabetes and at least one of the following conditions: dyslipidemia, hypertension, hyperuricemia, a pro-coagulant state, atherosclerosis or truncal obesity.

According to a further aspect the present invention additionally provides a method of elevating intracellular cGMP in a mammal in need thereof comprising administering to said mammal a PDE9 inhibitor, a prodrug thereof, a pharmaceutically acceptable salt of said PDE9 inhibitor or of said prodrug, or a pharmaceutical composition comprising a PDE9 inhibitor. It is particularly preferred that type 2 diabetes, insulin resistance syndrome or hypertension is treated thereby.

According to a further aspect the invention additionally provides a method of treating hypertension in a mammal comprising administering to said mammal a PDE9 inhibitor, a prodrug thereof or a pharmaceutically acceptable salt of said PDE9 inhibitor or of said prodrug. It is particularly preferred that said PDE9 inhibitor is 5-(3-chloro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one, a prodrug thereof or a pharmaceutically acceptable salt thereof or of said prodrug.

DETAILED DESCRIPTION OF THE INVENTION

The PDE9 inhibitors used in the pharmaceutical compositions and methods of this invention may be prepared as set forth in the Examples provided below or by following procedures analogous to those set forth in U.S. Patent No. 6,235,742 B1, which is incorporated herein by reference.

The pharmaceutically acceptable salts of the cGMP PDE9 inhibitor compounds as disclosed herein for use in the treatment of the insulin resistance syndrome in accordance with the present invention which contain a basic centre are, for example, non-toxic acid addition salts formed with inorganic acids such as hydrochloric, hydrobromic, hydroiodic, sulphuric and phosphoric acid, with carboxylic acids or with organo-sulphonic acids.

Examples include the HCl, HBr, HI, sulphate or bisulphate, nitrate, phosphate or hydrogen phosphate, acetate, benzoate, succinate, saccharate, fumarate, maleate, lactate, citrate, tartrate, gluconate, camsylate, methanesulphonate, ethanesulphonate, benzene-sulphonate, p-toluenesulphonate and pamoate salts. The cGMP PDE9 inhibitor
5 compounds for use in the present invention can also provide pharmaceutically acceptable metal salts, in particular non-toxic alkali and alkaline earth metal salts, with bases. Examples include the sodium, potassium, aluminium, calcium, magnesium, zinc and diethanolamine salts. For a review on suitable pharmaceutical salts see Berge *et al*, J. Pharm, Sci., 66, 1-19, 1977.

10 The cGMP PDE9 inhibitor compounds suitable for use in accordance with the present invention, their pharmaceutically acceptable salts, and pharmaceutically acceptable solvates of either entity can be administered alone but, in human therapy will generally be administered in admixture with a suitable pharmaceutical excipient diluent or carrier selected with regard to the intended route of administration and standard
15 pharmaceutical practice.

For example, the cGMP PDE9 inhibitor compounds suitable for use in accordance with the present invention or salts or solvates thereof can be administered orally, buccally or sublingually in the form of tablets, capsules (including soft gel capsules), multi-particulate, gels, films, ovules, elixirs, solutions or suspensions, which may contain
20 flavouring or colouring agents, for immediate-, delayed-, modified-, sustained-, dual-, controlled-release or pulsatile delivery applications. Such compounds may also be administered via fast dispersing or fast dissolving dosages forms or in the form of a high energy dispersion or as coated particles. Suitable pharmaceutical formulations may be in coated or un-coated form as desired.

25 Such solid pharmaceutical compositions, for example, tablets may contain excipients such as microcrystalline cellulose, lactose, sodium citrate, calcium carbonate, dibasic calcium phosphate, glycine and starch (preferably corn, potato or tapioca starch), disintegrants such as sodium starch glycollate, croscarmellose sodium and certain complex silicates, and granulation binders such as polyvinylpyrrolidone,
30 hydroxypropylmethyl cellulose (HPMC), hydroxypropylcellulose (HPC), hydroxypropyl methylcellulose acetate succinate (HPMCAS), sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, stearic acid, glyceryl behenate and talc may be included.

35 Solid compositions of a similar type may also be employed as fillers in gelatin capsules or HPMC capsules. Preferred excipients in this regard include lactose, starch, a cellulose, milk sugar or high molecular weight polyethylene glycols. For aqueous

suspensions and/or elixirs, the cGMP PDE9 inhibitor compounds may be combined with various sweetening or flavouring agents, colouring matter or dyes, with emulsifying and/or suspending agents and with diluents such as water, ethanol, propylene glycol and glycerin, and combinations thereof.

5 Modified release and pulsatile release dosage forms may contain excipients such as those detailed for immediate release dosage forms together with additional excipients that act as release rate modifiers, these being coated on and/or included in the body of the device. Release rate modifiers include, but are not exclusively limited to, HPMC, HPMCAS, methyl cellulose, sodium carboxymethylcellulose, ethyl cellulose, cellulose
10 acetate, polyethylene oxide, Xanthan gum, Carbomer, ammonio methacrylate copolymer, hydrogenated castor oil, carnauba wax, paraffin wax, cellulose acetate phthalate, hydroxypropylmethyl cellulose phthalate, methacrylic acid copolymer and mixtures thereof. Modified release and pulsatile release dosage forms may contain one or a combination of release rate modifying excipients. Release rate modifying excipients
15 maybe present both within the dosage form i.e. within the matrix, and/or on the dosage form, i.e., upon the surface or coating.

Fast dispersing or dissolving dosage formulations (FDDFs) may contain the following ingredients: aspartame, acesulfame potassium, citric acid, croscarmellose sodium, crospovidone, diascorbic acid, ethyl acrylate, ethyl cellulose, gelatin,
20 hydroxypropylmethyl cellulose, magnesium stearate, mannitol, methyl methacrylate, mint flavouring, polyethylene glycol, fumed silica, silicon dioxide, sodium starch glycolate, sodium stearyl fumarate, sorbitol, xylitol. The terms dispersing or dissolving as used herein to describe FDDFs are dependent upon the solubility of the drug substance used i.e. where the drug substance is insoluble a fast dispersing dosage form can be prepared
25 and where the drug substance is soluble a fast dissolving dosage form can be prepared.

The cGMP PDE9 inhibitor compounds suitable for use in accordance with the present invention can also be administered parenterally, for example, intracavernosally, intravenously, intra-arterially, intraperitoneally, intrathecally, intraventricularly, intraurethrally, intrasternally, intracranially, intramuscularly or subcutaneously, or they
30 may be administered by infusion or needle-free techniques. For such parenteral administration they are best used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or glucose to make the solution isotonic with blood. The aqueous solutions should be suitably buffered (preferably to a pH of from 3 to 9), if necessary. The preparation of suitable parenteral formulations
35 under sterile conditions is readily accomplished by standard pharmaceutical techniques well-known to those skilled in the art.

For oral and parenteral administration to human patients, the daily dosage level of the cGMP PDE9 inhibitor compounds for use in the present invention or salts or solvates thereof will usually be from 1 to 500 mg (in single or divided doses). A preferred dosage range is about 1 mg to about 100 mg. For the treatment of IRS the dosage may be via
5 single dose, divided daily dose, multiple daily dose, continuous (chronic) daily dosing for a specified period which may be from one to five or 5 or more, such as up to 10 or more days. Alternatively the treatment of IRS may be affected by continuous dosing, such as for example, via a controlled release dosage form wherein such continuous dosage form can be administered on a daily basis for a number of days or wherein such continuous
10 dosing can be affected via a slow-release formulation which doses for more than one day at a time.

Thus, for example, tablets or capsules of the cGMP PDE9 inhibitor compounds suitable for use in accordance with the present invention or salts or solvates thereof may contain from 1 mg to 250 mg of active compound for administration singly or two or more
15 at a time, as appropriate. Preferred tablets or capsules will contain about 1 mg to about 50 mg of active compound for administration singly or two or more at a time, as appropriate. The physician in any event will determine the actual dosage which will be most suitable for any individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case. There
20 can, of course, be individual instances where higher or lower dosage ranges are merited and such are within the scope of this invention.

The cGMP PDE9 inhibitor compounds suitable for use in accordance with the present invention can also be administered intranasally or by inhalation and are conveniently delivered in the form of a dry powder inhaler or an aerosol spray
25 presentation from a pressurised container, pump, spray or nebuliser with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, a hydrofluoroalkane such as 1,1,1,2-tetrafluoroethane (HFA 134A [trade mark] or 1,1,1,2,3,3,3-heptafluoropropane (HFA 227EA [trade mark]), carbon dioxide or other suitable gas. In the case of a pressurised aerosol, the dosage unit may
30 be determined by providing a valve to deliver a metered amount. The pressurised container, pump, spray or nebuliser may contain a solution or suspension of the active compound, e.g. using a mixture of ethanol and the propellant as the solvent, which may additionally contain a lubricant, e.g. sorbitan trioleate. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated to contain a
35 powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

Aerosol or dry powder formulations are preferably arranged so that each metered dose or "puff" contains from 1 to 50 mg of a compound of the invention for delivery to the patient. The overall daily dose with an aerosol will be in the range of from 1 to 50 mg which may be administered in a single dose or, more usually, in divided doses throughout the day.

The cGMP PDE9 inhibitor compounds suitable for use in accordance with the present invention may also be formulated for delivery via an atomiser. Formulations for atomiser devices may contain the following ingredients as solubilisers, emulsifiers or suspending agents: water, ethanol, glycerol, propylene glycol, low molecular weight polyethylene glycols, sodium chloride, fluorocarbons, polyethylene glycol ethers, sorbitan trioleate, oleic acid.

Alternatively, the cGMP PDE9 inhibitor compounds suitable for use in accordance with the present invention or salts or solvates thereof can be administered in the form of a suppository or pessary, or they may be applied topically in the form of a gel, hydrogel, lotion, solution, cream, ointment or dusting powder. The cGMP PDE9 inhibitor compounds suitable for use in accordance with the present invention or salts or solvates thereof may also be dermally or transdermally administered, for example, by the use of a skin patch. They may also be administered by the pulmonary or rectal routes.

The compounds may also be administered by the ocular route. For ophthalmic use, the compounds can be formulated as micronised suspensions in isotonic, pH adjusted, sterile saline, or, preferably, as solutions in isotonic, pH adjusted, sterile saline, optionally in combination with a preservative such as a benzylalkonium chloride. Alternatively, they may be formulated in an ointment such as petrolatum.

For application topically to the skin, the cGMP PDE9 inhibitor compounds suitable for use in accordance with the present invention or salts or solvates thereof can be formulated as a suitable ointment containing the active compound suspended or dissolved in, for example, a mixture with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, they can be formulated as a suitable lotion or cream, suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, a polyethylene glycol, liquid paraffin, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

The cGMP PDE9 inhibitor compounds suitable for use in accordance with the present invention may also be used in combination with a cyclodextrin. Cyclodextrins are known to form inclusion and non-inclusion complexes with drug molecules. Formation of a

drug-cyclodextrin complex may modify the solubility, dissolution rate, bioavailability and/or stability property of a drug molecule. Drug-cyclodextrin complexes are generally useful for most dosage forms and administration routes. As an alternative to direct complexation with the drug the cyclodextrin may be used as an auxiliary additive, e.g. as a carrier, diluent or solubiliser. Alpha-, beta- and gamma-cyclodextrins are most commonly used and suitable examples are described in International Patent Application Publication Nos. WO91/11172, WO94/02518 and WO98/55148.

Generally, in humans, oral administration is the preferred route, being the most convenient. In circumstances where the recipient suffers from a swallowing disorder or from impairment of drug absorption after oral administration, the drug may be administered parenterally, sublingually or buccally.

For veterinary use, a compound, or a veterinarily acceptable salt thereof, or a veterinarily acceptable solvate or pro-drug thereof, is administered as a suitably acceptable formulation in accordance with normal veterinary practice and the veterinary surgeon will determine the dosing regimen and route of administration which will be most appropriate for a particular animal.

It is to be appreciated that all references herein to treatment include curative, palliative and prophylactic treatment.

The present invention additionally comprises the use of a combination of a first and a second compound for the treatment of the insulin resistance syndrome or type 2 diabetes. The first compound of the combination is a cGMP PDE9 inhibitor compound as defined herein. The second compound of the combination is a naturally occurring or synthetic prostaglandin or ester thereof; an α -adrenergic receptor antagonist compound (also known as α -adrenoreceptors, α -receptors or α -blockers); a nitric oxide donor (also known as NO-donor or NO-agonist); a potassium channel opener or potassium channel modulator; a dopaminergic agent; a vasodilator agent; a thromboxane A2 agonist; an ergot alkaloid; a compound which modulates the action of a naturetic factor, particularly a compound which modulates the action of atrial naturetic factor (also known as atrial naturetic peptide), B type and C type naturetic factors; an angiotensin receptor antagonist; a substrate for NO-synthase; a calcium channel blocker; an antagonist of endothelin receptors; an inhibitor of endothelin converting enzyme; a cholesterol lowering agent such as an HMG-CoA reductase inhibitor; an antiplatelet or antithrombotic agent; an insulin sensitizing agent such as a glitazone; an insulin secretagogue such as a sulfonylurea; an acetylcholinesterase inhibitor; an estrogen receptor modulator; a PDE5 inhibitor; a PDE11 inhibitor; a neuropeptide Y (NPY) inhibitor, preferably an NPY5 inhibitor and even more preferably an NPY1 inhibitor, said NPY inhibitor having an IC50

of less than 100nM, and more preferably an IC₅₀ of less than 50nM; a vasoactive intestinal protein (VIP) or a VIP mimetic, more particularly a VIP which is mediated by one or more of the VIP receptor subtypes VPAC1, VPAC or PACAP (pituitary adenylate cyclase activating peptide); a VIP receptor agonist; a VIP analogue or fragment; an α -adrenoreceptor antagonist/VIP combination (e.g., Invicorp[®], Aviptadil); a serotonin receptor agonist, antagonist or modulator, more particularly, a modulator for 5HT1A; a testosterone replacement agent; estrogen; a combination of estrogen and medroxyprogesterone; a combination of estrogen and medroxyprogesterone acetate (MPA); a combination of estrogen and a methyl testosterone hormone replacement therapy agent (e.g., HRT); a modulator of transporters for noradrenaline, dopamine or serotonin; a purinergic receptor agonist or modulator; a neurokinin (NK) receptor antagonist; an opioid receptor agonist, antagonist or modulator, preferably an agonist for the ORL-1 receptor; an oxytocin/vasopressin receptor modulator or agonist, preferably a selective oxytocin agonist or modulator; a cannabinoid receptor modulator; a central nervous system (CNS) active agent; an angiotensin-converting enzyme inhibitor; a combination of an angiotensin converting-enzyme inhibitor and a neutral endopeptidase; L-Dopa; a combination of L-Dopa and carbidopa; a steroidal anti-inflammatory agent; a non-steroidal anti-inflammatory agent; a protein kinase C- β inhibitor; an AMP-activated protein kinase activator; insulin; a weight loss agent; a dipeptidyl peptidase IV (DPP IV) inhibitor; a glucagon antagonist; an I kappa B kinase- β (IKK- β) inhibitor such as salicylate; a PTP1B inhibitor; an agent that reduces the levels of PTP1B levels using antisense technology; a glycogen synthase kinase-3 inhibitor; a GLP-1 agonist; a PPAR- γ agonist; a PPAR- γ antagonist; a PPAR- α agonist; a dual PPAR- α /PPAR- γ agonist; a RXR antagonist; a biguanide such as metformin, a glycogen phosphorylase inhibitor; a sorbitol dehydrogenase inhibitor (SDI); an aldose reductase inhibitor (ARI); a soluble guanylate cyclase (sGC) activator; growth hormone; or a growth hormone secretagogue.

Any naturally occurring or synthetic prostaglandin or ester thereof may be used as the second compound of a combination of this invention. Suitable prostaglandins for use herein include alprostadil, prostaglandin E₁, prostaglandin E₂, 13, 14 - dihydroprostaglandin E₁, prostaglandin E₂, eprostinol, natural synthetic and semi-synthetic prostaglandins and derivatives thereof including those described in International Patent Application Publication No. WO00/33825 and U.S. Patent No. 6,037,346; PGE₀, PGE₁, PGA₁, PGB₁, PGF₁ α , 19-hydroxy PGA₁, 19-hydroxy - PGB₁, PGE₂, PGB₂, 19-hydroxy-PGA₂, 19-hydroxy-PGB₂, PGE₃ α , carboprost tromethamine dinoprost, tromethamine, dinoprostone, lipoprost, gemeprost, metenoprost, sulprostone, tiaprost and moxisylyate.

The disclosures made in U.S. patents, International patent applications and all other references mentioned herein are hereby incorporated by reference.

Any α -adrenergic receptor antagonist compound may be used as the second compound of a combination of this invention. Suitable α -adrenergic receptor antagonists for use herein include the α -adrenergic receptor blockers described in International Patent Application Publication No. WO99/30697. Selective α_1 -adrenoceptor, α_2 -adrenoceptor blockers and non-selective adrenoceptor blockers may also be used as the second α -adrenergic receptor antagonist compound of this invention. Suitable α_1 -adrenoceptor blockers include phentolamine, phentolamine mesylate, trazodone, alfuzosin, indoramin, naftopidil, tamsulosin, dapiprazole, phenoxybenzamine, idazoxan, efaraxan, yohimbine, rauwolfia alkaloids, Recordati 15/2739, SNAP 1069, SNAP 5089, RS17053, SL 89.0591, doxazosin, terazosin, abanoquil and prazosin. Suitable α_2 -adrenoceptor blockers include those disclosed in U.S. Patent No. 6,037,346, dibenarnine, tolazoline, trimazosin and dibenamine. Suitable α -adrenergic receptors for use as the second compound of a combination of this invention are also described in U.S. Patent Nos. 4,188,390; 4,026,894; 3,511,836; 4,315,007; 3,527,761; 3,997,666; 2,503,059; 4,703,063; 3,381,009; 4,252,721 and 2,599,000. Other suitable α_2 -adrenoceptor blockers include clonidine, papaverine, papaverine hydrochloride, each of which may optionally be administered in the presence of a cariogenic agent such as, but not limited to, piraxamine.

Any nitric oxide donor (NO-donor or NO-agonist) compound may be used as the second compound of a combination of this invention. Suitable NO-donor compounds for use herein include organic nitrates, such as mono-, di- or tri-nitrates; organic nitrate esters such as glyceryl binitrate (also known as nitroglycerin), isosorbide 5-mononitrate, isosorbide dinitrate, pentaerythritol tetranitrate, erythrityl tetranitrate, amylnitrate, a diazenium diolate (NONOate), and 1,5-pentanedinitrate; sodium nitroprusside (SNP); 3-morpholinosydnonimine molsidomine; S-nitroso- N-acetyl penicilliamine (SNAP); S-nitroso-N-glutathione (SNO-GLU); N-hydroxy-L-arginine; linsidomine; linsidomine chlorohydrate; (SIN-1) S-nitroso-N-cysteine; L-arginine; ginseng; zizphi fructus; molsidomine; Re - 2047; and nitrosylated maxisylate derivatives such as NMI-678-11 and NMI-937 (International Patent Application Publication No. WO00/12075).

Any potassium channel opener or modulator may be used as the second compound of a combination of this invention. Suitable potassium channel openers/modulators for use herein include nicorandil, cromokalim, levromokalim, lemakalim, pinacidil, cliazoxide, minoxidil, charybdotoxin, glyburide, glipizide, 4-aminopyridine and barium chloride (BaCl_2).

Any dopaminergic agent may be used as the second compound of a combination of this invention. Preferred dopaminergic agents include apomorphine and selective D₂, D₃ and D₂/D₃ agonists such as pramipexole, ropirinol (International Patent Application Publication No. WO00/23056), L-Dopa, L-Dopa in combination with carbidopa, PNU95666 (International Patent Application Publication No. WO00/40226).

Any vasodilator agent may be used as the second compound of a combination of this invention. Suitable vasodilator agents for use herein include nimodipine, pinacidil, cyclandelate, isoxsuprine, chloroprumazine, haloperidol, Rec 15/2739 and trazodone.

Any ergot alkaloid may be used as the second compound of a combination of this invention. Suitable ergot alkaloids include those disclosed in U.S. Patent No. 6,037,346; acetergamine, brazergoline, bromerguride, cianergoline, delorgotril, disulergine, ergonovine maleate, ergotamine tartrate, etisulergine, lergotril, lysergide, mesulergine, metergoline, metergotamine, nicergoline, pergolide, propisergide, proterguride, terguride.

Any angiotensin receptor antagonist may be used as the second compound of a combination of this invention. Suitable angiotensin receptor antagonists include losartan, candersartan, eprosartan, irbesartan and valsartan.

Any substrate for NO-synthase may be used as the second compound of a combination of this invention. Suitable NO-synthase substrates include, *inter alia*, L-arginine.

Any calcium channel blocker may be used as the second compound of a combination of this invention. Suitable calcium channel blockers include, amlodipine (amlodipine besylate is also known as Norvasc®), bepridil, which may be prepared as disclosed in U.S. Patent No. 3,962, 238 or U.S. Reissue No. 30,577; clentiazem, which may be prepared as disclosed in U.S. Patent No. 4,567,175; diltiazem, which may be prepared as disclosed in U.S. Patent No. 3,562, fendiline, which may be prepared as disclosed in U.S. Patent No. 3,262,977; gallopamil, which may be prepared as disclosed in U.S. Patent No. 3,261,859; mibefradil, which may be prepared as disclosed in U.S. Patent No. 4,808,605; prenylamine, which may be prepared as disclosed in U.S. Patent No. 3,152,173; semotiadil, which may be prepared as disclosed in U.S. Patent No. 4,786,635; terodiline, which may be prepared as disclosed in U.S. Patent No. 3,371,014; verapamil, which may be prepared as disclosed in U.S. Patent No. 3,261,859; aranipine, which may be prepared as disclosed in U.S. Patent No. 4,572,909; barnidipine, which may be prepared as disclosed in U.S. Patent No. 4,220,649; benidipine, which may be prepared as disclosed in European Patent Application Publication No. 106,275; cilnidipine, which may be prepared as disclosed in U.S. Patent No. 4,672,068; efonidipine, which may be prepared as disclosed in U.S. Patent No. 4,885,284; elgodipine, which may be prepared as disclosed in U.S. Patent No.

4,952,592; felodipine, which may be prepared as disclosed in U.S. Patent No. 4,264,611; isradipine, which may be prepared as disclosed in U.S. Patent No. 4,466,972; lacidipine, which may be prepared as disclosed in U.S. Patent No. 4,801,599; lercanidipine, which may be prepared as disclosed in U.S. Patent No. 4,705,797; manidipine, which may be prepared as disclosed in U.S. Patent No. 4,892,875; nicardipine, which may be prepared as disclosed in U.S. Patent No. 3,985,758; nifedipine, which may be prepared as disclosed in U.S. Patent No. 3,485,847; nilvadipine, which may be prepared as disclosed in U.S. Patent No. 4,338,322; nimodipine, which may be prepared as disclosed in U.S. Patent No. 3,799,934; nisoldipine, which may be prepared as disclosed in U.S. Patent No. 4,154,839; nitrendipine, which may be prepared as disclosed in U.S. Patent No. 3,799,934; cinnarizine, which may be prepared as disclosed in U.S. Patent No. 2,882,271; flunarizine, which may be prepared as disclosed in U.S. Patent No. 3,773,939; lidoflazine, which may be prepared as disclosed in U.S. Patent No. 3,267,104; lomerizine, which may be prepared as disclosed in U.S. Patent No. 4,663,325; bencyclane, which may be prepared as disclosed in Hungarian Patent No. 151,865; etafenone, which may be prepared as disclosed in German Patent No. 1,265,758; and perhexiline, which may be prepared as disclosed in British Patent No. 1,025,578.

Any one cholesterol lowering agent may be used as the second compound of a combination of this invention. Suitable cholesterol lowering agents include vastatins such as simvastatin, disclosed in U.S. 4,444,784; pravastatin, disclosed in U.S. 4,346,227; cerivastatin, disclosed in U.S. 5,502,199; mevastatin, disclosed in U.S. 3,983,140; velostatin, disclosed in U.S. 4,448,784 and U.S. 4,450,171; fluvastatin, disclosed in U.S. 4,739,073; compactin, disclosed in U.S. 4,804,770; lovastatin, disclosed in U.S. 4,231,938; dalvastatin, disclosed in European Patent Application Publication No. 738510 A2; fluindostatin, disclosed in European Patent Application Publication No. 363934 A1; atorvastatin, disclosed in U.S. Patent No. 4,681,893; atorvastatin calcium (atorvastatin calcium is also known as Lipitor®), disclosed in U.S. Patent No. 5,273,995; and dihydrocompactin, disclosed in U.S. 4,450,171. Other suitable cholesterol lowering agents include fibrates.

Any antiplatelet and antithrombotic agent may be used as the second compound of a combination of this invention. Suitable antiplatelet and antithrombotic agents include, e.g., tPA, uPA, warfarin, hirudin and other thrombin inhibitors, heparin and thromboplastin activating factor inhibitors.

Any insulin sensitising agent may be used as the second compound of a combination of this invention. Suitable insulin sensitizing agents include Avandia®, Actos® and hypoglycaemic agents such as, but not limited to, sulfonylureas such as glipizide,

metformin and acarbose.

Any acetylcholinesterase inhibitor may be used as the second compound of a combination of this invention. A suitable acetylcholinesterase inhibitor is, e.g., donepezil.

Any estrogen receptor modulator, estrogen agonist or estrogen antagonist may be used as the second compound of a combination of this invention. Suitable estrogen receptor modulators, estrogen agonists or estrogen antagonists include the compounds disclosed in International Patent Application Publication No. WO96/21656 and U.S. Patent No. 5,552,412. Preferred such compounds include raloxifene, lasofoxifene, (-)-cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydronaphthalene-2-ol and pharmaceutically acceptable salts thereof.

Any PDE5 or PDE11 inhibitor may be used as the second compound of a combination of this invention. It is particularly preferred that a PDE5 inhibitor be used as the second compound of this invention. Suitable PDE5 inhibitors include the pyrazolo [4,3-d]pyrimidin-7-ones disclosed in EP-A-0463756; the pyrazolo [4,3-d]pyrimidin-7-ones disclosed in EP-A-0526004; the pyrazolo [4,3-d]pyrimidin-7-ones disclosed in published international patent application WO 93/06104; the isomeric pyrazolo [3,4-d]pyrimidin-4-ones disclosed in International Patent Application Publication No. WO93/07149; the quinazolin-4-ones disclosed in International Patent Application Publication No. WO93/12095; the pyrido [3,2-d]pyrimidin-4-ones disclosed in International Patent Application Publication No. WO94/05661; the purin-6-ones disclosed in International Patent Application Publication No. WO94/00453; the pyrazolo [4,3-d]pyrimidin-7-ones disclosed in International Patent Application Publication No. WO98/49166; the pyrazolo [4,3-d]pyrimidin-7-ones disclosed in International Patent Application Publication No. WO99/54333; the pyrazolo [4,3-d]pyrimidin-4-ones disclosed in EP-A-0995751; the pyrazolo [4,3-d]pyrimidin-7-ones disclosed in International Patent Application Publication No. WO00/24745; the pyrazolo [4,3-d]pyrimidin-4-ones disclosed in EP-A-0995750; the compounds disclosed in International Patent Application Publication No. WO95/19978; the compounds disclosed in International Patent Application Publication No. WO99/24433; the pyrazolo [4,3-d]pyrimidin-7-ones disclosed in International Patent Application Publication No. WO01/27112; the pyrazolo [4,3-d]pyrimidin-7-ones disclosed in International Patent Application Publication No. WO01/27113; the compounds disclosed in EP-A-1092718; the compounds disclosed in EP-A-1092719; and the compounds disclosed in International Patent Application Publication No. WO93/07124.

Preferred PDE5 inhibitors for use as a second compound in a combination of this invention include: 5-[2-ethoxy-5-(4-methyl-1-piperazinylsulphonyl)phenyl]-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (sildenafil) also known as 1-[[3-(6,7-

dihydro-1-methyl-7-oxo-3-propyl-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-4-ethoxyphenyl)sulphonyl]-4-methylpiperazine (see EP-A-0463756); 5-(2-ethoxy-5-morpholinoacetylphenyl)-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (see EP-A-0526004); 3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-n-propoxyphenyl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (see WO98/49166); 3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-(2-methoxyethoxy)pyridin-3-yl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (see WO99/54333); 6-benzo[1,3]dioxol-5-yl-2-methyl-2,3,6,7,12,12a-hexahydro-pyrazino[1',2':1,6]pyrido[3,4-b]indole-1,4-dione (cialis); (+)-3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-(2-methoxy-1(R)-methylethoxy)pyridin-3-yl]-2-methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one, also known as 3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-((1R)-2-methoxy-1-methylethyl)oxy]pyridin-3-yl]-2-methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (see WO99/54333); 5-[2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-2-[2-methoxyethyl]-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one, also known as 1-[6-ethoxy-5-[3-ethyl-6,7-dihydro-2-(2-methoxyethyl)-7-oxo-2H-pyrazolo[4,3-d]pyrimidin-5-yl]-3-pyridylsulphonyl]-4-ethylpiperazine (see WO01/27113, Example 8); 5-[2-iso-butoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-2-(1-methylpiperidin-4-yl)-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (see WO01/27113, Example 15); 5-[2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-2-phenyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (see WO01/27113, Example 66); 5-(5-acetyl-2-propoxy-3-pyridinyl)-3-ethyl-2-(1-isopropyl-3-azetidiny)-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (see WO01/27112, Example 124); 5-(5-acetyl-2-butoxy-3-pyridinyl)-3-ethyl-2-(1-ethyl-3-azetidiny)-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (see WO01/27112, Example 132); (6R,12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione (IC-351), i.e. the compound of examples 78 and 95 of published international application WO95/19978, as well as the compound of examples 1, 3, 7 and 8; 2-[2-ethoxy-5-(4-ethylpiperazin-1-yl-1-sulphonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-f][1,2,4]triazin-4-one (vardenafil) also known as 1-[[3-(3,4-dihydro-5-methyl-4-oxo-7-propylimidazo[5,1-f]-as-triazin-2-yl)-4-ethoxyphenyl)sulphonyl]-4-ethylpiperazine, i.e. the compound of examples 20, 19, 337 and 336 of published international application WO99/24433; the compound of example 11 in WO93/07124 (EISA); and compounds 3 and 14 from Rotella D P, *J. Med. Chem.*, 2000, 43, 1257.

Still other type cGMP PDE5 inhibitors useful in conjunction with the present invention include: 4-bromo-5-(pyridylmethylamino)-6-[3-(4-chlorophenyl)-propoxy]-3(2H)pyridazinone; 1-[4-[(1,3-benzodioxol-5-ylmethyl)amino]-6-chloro-2-quinazoliny]-4-

piperidine-carboxylic acid, monosodium salt; (+)-cis-5,6a,7,9,9a-hexahydro-2-[4-(trifluoromethyl)-phenylmethyl-5-methyl-cyclopent-4,5]imidazo[2,1-b]purin-4(3H)one; furazlocillin; cis-2-hexyl-5-methyl-3,4,5,6a,7,8,9,9a-octahydrocyclopent[4,5]imidazo[2,1-b]purin-4-one; 3-acetyl-1-(2-chlorobenzyl)-2-propylindole-6-carboxylate; 3-acetyl-1-(2-chlorobenzyl)-2-propylindole-6-carboxylate; 4-bromo-5-(3-pyridylmethylamino)-6-(3-(4-chlorophenyl) propoxy)-3-(2H)pyridazinone; 1-methyl-5-(5-morpholinoacetyl-2-n-propoxyphenyl)-3-n-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one; 1-[4-[(1,3-benzodioxol-5-ylmethyl)amino]-6-chloro-2-quinazoliny]-4-piperidinecarboxylic acid, monosodium salt; Pharmaprojects No. 4516 (Glaxo Wellcome); Pharmaprojects No. 5051 (Bayer); Pharmaprojects No. 5064 (Kyowa Hakko; see WO 96/26940); Pharmaprojects No. 5069 (Schering Plough); GF-196960 (Glaxo Wellcome); E-8010 and E-4010 (Eisai); Bay-38-3045 & 38-9456 (Bayer) and Sch-51866; selected from: sildenafil, 5-(2-ethoxy-5-morpholinoacetylphenyl)-1-ethyl-3-n-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one, 3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-n-propoxyphenyl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one, 3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-(2-methoxyethoxy)pyridin-3-yl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one, 5-[2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-2-(2-methoxyethyl)-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one and 1-[[3-(3,4-dihydro-5-methyl-4-oxo-7-propylimidazo[5,1-f]as-triazin-2-yl)-4-ethoxyphenyl]sulphonyl]-4-ethylpiperazine or a pharmaceutically acceptable salt, solvate, pro-drug or polymorph thereof.

More preferred cGMP PDE5 inhibitors for use as the second compound in a combination of this invention include sildenafil, sildenafil citrate (also known as Viagra®; 5-(5-acetyl-2-butoxy-3-pyridinyl)-3-ethyl-2-(1-ethyl-3-azetidiny)-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one; 2-[2-ethoxy-5-(4-ethylpiperazin-1-yl-1-sulphonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-f][1,2,4]triazin-4-one (vardenafil); 6-benzo[1,3]dioxol-5-yl-2-methyl-2,3,6,7,12,12a-hexahydro-pyrazino[1',2':1,6]pyrido[3,4-b]indole-1,4-dione (cialis); and 5-[2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-1-(2-methoxyethyl)-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one.

Any melanocortin receptor agonist, melanocortin receptor modulator or melanocortin receptor enhancer may be used as the second compound of a combination of this invention. Suitable melanocortin receptor agonists, modulators or enhancers include melanotan II; PT-14; PT-141; and compounds disclosed in International Patent Application Publication Nos. WO99/64002, WO00/74679, WO99/55679, WO01/05401, WO00/58361, WO01/14879, WO01/13112 and WO99/54358.

Any serotonin receptor agonist, antagonist or modulator may be used as the

second compound in a combination of this invention. It is particularly preferred to use agonists, antagonists or modulators of 5HT1A. Suitable such agonists, antagonists or modulators include VML 670; 5HT2A; 5HT2C; 5HT3; and 5HT6 receptors, including those described in International Patent Application Publication Nos. WO99/02159, WO00/02550 and WO00/28993.

Any testosterone replacement agent may be used as the second compound in a combination of this invention. Suitable testosterone replacement agents include dehydroandrostendione, testosterone (Tostrelle), dihydrotestosterone and testosterone implants.

Any hormone replacement therapy (HRT) agent may be used as the second compound of a combination of this invention. Suitable HRT agents include Premarin[®], Cenestin[®], Oestrofeminal[®], Equin[®], Estrace[®], Estrofem[®], Elleste Solo[®], Estring[®], Estraderm TTS[®], Estraderm Matrix[®], Dermestril[®], Premphase[®], Preempro[®], Prempak[®], Premique[®], Estratest[®], Estratest HS[®] and Livial[®] (tibolone).

Any modulator of transporters for noradrenaline, dopamine and/or serotonin may be used as the second compound of a combination of this invention. Suitable such modulators include bupropion and GW-320659.

Any neurokinin (NK) receptor antagonist may be used as the second compound of a combination of this invention. Suitable NK receptor antagonists include those described in International Patent Application Publication No. WO99/64008.

Any angiotensin converting enzyme inhibitor (ACE inhibitor) may be used as the second compound of a combination of this invention. Suitable ACE inhibitors include, but are not limited to: alacepril, which may be prepared as disclosed in U.S. Patent No. 4,248,883; benazepril, which may be prepared as disclosed in U.S. Patent No. 4,410,520; captopril, which may be prepared as disclosed in U.S. Patent Nos. 4,046,889 and 4,105,776; ceronapril, which may be prepared as disclosed in U.S. Patent No. 4,452,790; delapril, which may be prepared as disclosed in U.S. Patent No. 4,385,051; enalapril, which may be prepared as disclosed in U.S. Patent No. 4,374,829; fosinopril, which may be prepared as disclosed in U.S. Patent No. 4,337,201; imadapril, which may be prepared as disclosed in U.S. Patent No. 4,508,727; lisinopril, which may be prepared as disclosed in U.S. Patent No. 4,555,502; moveltopril, which may be prepared as disclosed in Belgian Patent No. 893,553; perindopril, which may be prepared as disclosed in U.S. Patent No. 4,508,729; quinapril, which may be prepared as disclosed in U.S. Patent No. 4,344,949; ramipril, which may be prepared as disclosed in U.S. Patent No. 4,587,258; spirapril, which may be prepared as disclosed in U.S. Patent No. 4,470,972; temocapril, which may be prepared as disclosed in U.S. Patent No. 4,699,905; and trandolapril, which may be

prepared as disclosed in U.S. Patent No. 4,933,361.

Any compound which is a combined inhibitor of angiotensin-converting enzyme and neutral endopeptidase may be used as the second compound of a combination of this invention. A suitable such combined inhibitor is, e.g., omapatrilat.

5 Any protein kinase C- β inhibitor may be used as the second compound of a combination of this invention. A suitable protein kinase C- β inhibitor is, e.g., LY333531.

Any activator of AMP-activated protein kinase may be used as the second compound of a combination of this invention. A suitable such activator is, e.g., 5-amino-4-imidazolecarboxamide ribonucleoside.

10 Any weight loss agent may be used as the second compound of a combination of this invention. Suitable weight loss agents include sibutramine and orlistat.

Any dipeptidyl peptidase IV (DPPIV) inhibitor may be used as the second compound of a combination of this invention. Suitable DPPIV inhibitors include NVP DPP728 and P32/98.

15 Any glucagon antagonist may be used as the second compound of a combination of this invention. A suitable glucagon antagonist is, e.g., NNC25-2504.

Any IKK- β inhibitor may be used as the second compound of a combination of this invention. A suitable IKK- β inhibitor is, e.g., salicylate.

Any PTP1B inhibitor may be used as the second compound of a combination of this invention. A suitable PTP1B inhibitor is, e.g., PTP112.

20 Any glycogen synthase kinase-3 (GSK-3) inhibitor may be used as the second compound of a combination of this invention. A suitable GSK-3 inhibitor is, e.g., Chir98014

Any GLP-1 agonist may be used as the second compound of a combination of this invention. Suitable GLP-1 agonists include GLP1, NN-2211 and exendin 4.

Any PPAR- γ agonist may be used as the second compound of a combination of this invention. Suitable PPAR- γ agonists include Rezulin[®], Avandia[®], Actos[®] or CS011.

Any PPAR- γ antagonist may be used as the second compound of a combination of this invention. A suitable PPAR- γ antagonist is, e.g., bisphenol A diglycidyl ether (BADGE).

30 Any PPAR- α agonist may be used as the second compound of a combination of this invention. A suitable PPAR- α agonist is, e.g., fenofibrate.

Any dual PPAR- α /PPAR- γ agonist may be used as the second compound of a combination of this invention. Suitable such dual agonists include farglitazar, GW1929, DRF2725, AZ242 and KRP 297.

Any RXR antagonist may be used as the second compound of a combination of this invention. A suitable RXR antagonist is, e.g., HX531.


Any glycogen phosphorylase inhibitor may be used as the second compound of a combination of this invention. A suitable glycogen phosphorylase inhibitor is, e.g., CP-
5 316819.

Any sorbitol dehydrogenase inhibitor (SDI) may be used as the second compound of a combination of this invention. Suitable SDIs include those disclosed in International Patent Application Publication No. WO00/59510. A particularly preferred SDI is 1R-(4-(4-(4,6-dimethyl)-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl)-pyrimidin-2-yl)-ethanol.

10 Any aldose reductase inhibitor (ARI) may be used as the second compound of a combination of this invention. Suitable ARIs include zopolrestat, epalrestat, ponalrestat, zenarestat or fidarestat.

Other suitable ARIs for use as the second compound in a combination of this invention include compounds of the Formula ARI

15



ARI

prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein:

20 A is S, SO or SO₂;

R¹ and R² are each independently hydrogen or methyl;

R³ is Het¹, -CHR⁴Het¹ or NR⁶R⁷;

R⁴ is hydrogen or (C₁-C₃)alkyl;

R⁶ is (C₁-C₆)alkyl, aryl or Het²;

25 R⁷ is Het³;

Het¹ is pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, quinolyl, isoquinolyl, quinazolyl, quinoxalyl, phthalazinyl, cinnolyl, naphthyridinyl, pteridinyl, pyrazinopyrazinyl, pyrazinopyridazinyl, pyrimidopyridazinyl, pyrimidopyrimidyl, pyridopyrimidyl, pyridopyrazinyl, pyridopyridazinyl, pyrrolyl, furanyl, thienyl, imidazolyl, oxazolyl, thiazolyl, 30 pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzoxazolyl, benzothiazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, pyrrolopyridyl, furopyridyl, thienopyridyl, imidazolopyridyl, oxazolopyridyl, thiazolopyridyl, pyrazolopyridyl, isoxazolopyridyl, isothiazolopyridyl,

- pyrrolopyrimidyl, furopyrimidyl, thienopyrimidyl, imidazolopyrimidyl, oxazolopyrimidyl, thiazolopyrimidyl, pyrazolopyrimidyl, isoxazolopyrimidyl, isothiazolopyrimidyl, pyrrolopyrazinyl, furopyrazinyl, thienopyrazinyl, imidazolopyrazinyl, oxazolopyrazinyl, thiazolopyrazinyl, pyrazolopyrazinyl, isoxazolopyrazinyl, isothiazolopyrazinyl,
- 5 pyrrolopyridazinyl, furopyridazinyl, thienopyridazinyl, imidazolopyridazinyl, oxazolopyridazinyl, thiazolopyridazinyl, pyrazolopyridazinyl, isoxazolopyridazinyl or isothiazolopyridazinyl; Het¹ is optionally substituted with up to a total of four substituents each independently selected from halo, formyl, (C₁-C₆)alkoxycarbonyl, (C₁-C₆)alkylenyloxycarbonyl, (C₁-C₄)alkoxy-(C₁-C₄)alkyl, C(OH)R¹²R¹³, (C₁-C₄)alkylcarbonylamido, (C₃-C₇)cycloalkylcarbonylamido, phenylcarbonylamido, benzyl,
- 10 phenyl, naphthyl, imidazolyl, pyridyl, triazolyl, benzimidazolyl, oxazolyl, isoxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, thienyl, benzothiazolyl, pyrrolyl, pyrazolyl, quinolyl, isoquinolyl, benzoxazolyl, pyridazinyl, pyridyloxy, pyridylsulfonyl, furanyl, phenoxy, thiophenoxy, (C₁-C₄)alkylsulfenyl, (C₁-C₄)alkylsulfonyl, (C₃-C₇)cycloalkyl, (C₁-C₆)alkyl optionally substituted with up to three fluoro, or (C₁-C₄)alkoxy optionally substituted with up to five fluoro; said benzyl, phenyl, naphthyl, imidazolyl, pyridyl,
- 15 triazolyl, benzimidazolyl, oxazolyl, isoxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, thienyl, benzothiazolyl, pyrrolyl, pyrazolyl, quinolyl, isoquinolyl, benzoxazolyl, pyridazinyl, pyridyloxy, pyridylsulfonyl, furanyl, phenoxy, thiophenoxy, in the definition of substituents for Het¹ are optionally substituted with up to three substituents independently selected from hydroxy, halo, hydroxy-(C₁-C₄)alkyl, (C₁-C₄)alkoxy-(C₁-C₄)alkyl, (C₁-C₆)alkylsulfenyl, (C₁-C₆)alkylsulfinyl, (C₁-C₆)alkylsulfonyl, (C₁-C₆)alkyl optionally substituted with up to five fluoro and (C₁-C₆)alkoxy optionally substituted with up to five fluoro; said imidazolyl, oxazolyl, isoxazolyl, thiazolyl and pyrazolyl in the definition of substituents for Het¹ are
- 20 optionally substituted with up to two substituents independently selected from hydroxy, halo, C₁-C₆alkyl, hydroxy-(C₁-C₄)alkyl, (C₁-C₄)alkoxy-(C₁-C₄)alkyl, C₁-C₄alkyl-phenyl optionally substituted in the phenyl portion with one Cl, Br, OMe, Me or SO₂-phenyl wherein said SO₂-phenyl is optionally substituted in the phenyl portion with one Cl, Br, OMe, Me, (C₁-C₄)alkyl optionally substituted with up to five fluoro, or (C₁-C₄)alkoxy optionally substituted with up to three fluoro;
- 30 R¹² and R¹³ are each independently hydrogen or (C₁-C₄)alkyl;
- Het² and Het³ are each independently imidazolyl, pyridyl, triazolyl, benzimidazolyl, oxazolyl, isoxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, thienyl, benzothiazolyl, pyrrolyl, pyrazolyl, quinolyl, isoquinolyl, benzoxazolyl, pyridazinyl, pyridyloxy,
- 35 pyridylsulfonyl, furanyl, phenoxy, thiophenoxy; Het² and Het³ are each independently optionally substituted with up to a total of four substituents each independently selected

from halo, formyl, (C₁-C₆)alkoxycarbonyl, (C₁-C₆)alkylenyloxycarbonyl, (C₁-C₄)alkoxy-(C₁-C₄)alkyl, C(OH)R¹⁸R¹⁹, (C₁-C₄)alkylcarbonylamido, (C₃-C₇)cycloalkylcarbonylamido, phenylcarbonylamido, phenyl, naphthyl, imidazolyl, pyridyl, triazolyl, benzimidazolyl, oxazolyl, isoxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, thienyl, benzothiazolyl, 5 pyrrolyl, pyrazolyl, quinolyl, isoquinolyl, benzoxazolyl, pyridazinyl, pyridyloxy, pyridylsulfonyl, furanyl, phenoxy, thiophenoxy, (C₁-C₄)alkylsulfenyl, (C₁-C₄)alkylsulfonyl, (C₃-C₇)cycloalkyl, (C₁-C₄)alkyl optionally substituted with up to three fluoro or (C₁-C₄)alkoxy optionally substituted with up to five fluoro; said phenyl, naphthyl, imidazolyl, pyridyl, triazolyl, benzimidazolyl, oxazolyl, isoxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, 10 tetrazolyl, thienyl, benzothiazolyl, pyrrolyl, pyrazolyl, quinolyl, isoquinolyl, benzoxazolyl, pyridazinyl, pyridyloxy, pyridylsulfonyl, furanyl, phenoxy, thiophenoxy, in the definition of substituents for Het² and Het³ are optionally substituted with up to three substituents independently selected from hydroxy, halo, hydroxy-(C₁-C₄)alkyl, (C₁-C₄)alkoxy-(C₁-C₄)alkyl, (C₁-C₄)alkyl optionally substituted with up to five fluoro and (C₁-C₄)alkoxy 15 optionally substituted with up to five fluoro; said imidazolyl, oxazolyl, isoxazolyl, thiazolyl and pyrazolyl in the definition of substituents for Het² and Het³ are optionally substituted with up to two substituents independently selected from hydroxy, halo, hydroxy-(C₁-C₄)alkyl, (C₁-C₄)alkoxy-(C₁-C₄)alkyl, (C₁-C₄)alkyl optionally substituted with up to five fluoro and (C₁-C₄)alkoxy optionally substituted with up to three fluoro; and 20 R¹⁸ and R¹⁹ are each independently hydrogen or (C₁-C₄)alkyl, provided that when R³ is NR⁶R⁷, then A is SO₂. A particularly preferred compound of the Formula ARI for use as the second compound of a combination of this invention is 6-(5-chloro-3-methyl-benzofuran-2-sulfonyl)-2H-pyridazin-3-one.

Any soluble guanylate cyclase (sGC) activator may be used as the second 25 compound of a combination of this invention. Suitable sGC activators include BAY 41-2272 and BAY 41-8543.

Any growth hormone secretagogue may be used as the second compound of a combination of this invention. Suitable growth hormone secretagogues include those disclosed in U.S. Patent Nos. 6,124,264; 6,110,932; 6,278,000; and 6,251,902. A 30 particularly preferred growth hormone secretagogue is 2-amino-N-(2-(3a(R)-benzyl-2-methyl-3-oxo-2,3,3a,4,6,7-hexahydro-pyrazolo-[4,3c]pyridin-5-yl)-1(R)-benzyloxymethyl-2-oxo-ethyl)-isobutyramide.

Particularly preferred compounds for use as the second compound in the combinations and pharmaceutical compositions for use according to the present invention 35 include compounds selected from the following classes of compounds: insulin sensitizing agents, PDE5 inhibitors, protein kinase C-β inhibitors, AMP-activated protein kinase

activators, insulin, weight loss agents, PPAR- γ agonists, PPAR- α agonists, dual PPAR- γ /PPAR- α agonists, sorbitol dehydrogenase inhibitors and aldose reductase inhibitors, each as described above.

PDE9 inhibitor – TEST METHODS

5 Phosphodiesterase (PDE) inhibitory activity

Preferred PDE compounds suitable for use in accordance with the present invention are potent and cGMP PDE9 inhibitors. *In vitro* PDE inhibitory activities against cyclic guanosine 3',5'-monophosphate (cGMP) and cyclic adenosine 3',5'-monophosphate (cAMP) phosphodiesterases are determined by measurement of their IC₅₀ values (the
10 concentration of compound required for 50% inhibition of enzyme activity).

Phosphodiesterase 9 can be generated from full length human recombinant clones transfected into SF9 cells as described in Fisher et al., Journal of Biological Chemistry, 1998, 273, 15559 - 15564.

Assays are performed either using a modification of the "batch" method of W.J. Thompson *et al.* (Biochem., 1979, 18, 5228) or using a scintillation proximity assay for the
15 direct detection of AMP/GMP using a modification of the protocol described by Amersham plc under product code TRKQ7090/7100. In summary, the effect of PDE9 inhibitors is investigated by assaying a fixed amount of enzyme in the presence of varying inhibitor concentrations and low substrate, (cGMP in a 3:1 ratio unlabelled to [³H]-labeled at a
20 concentration of about 1/3 K_m) such that IC₅₀ \cong K_i. The final assay volume is made up to 100 μ l with assay buffer [20 mM Tris-HCl pH 7.4, 5 mM MgCl₂, 1 mg/ml bovine serum albumin]. Reactions are initiated with enzyme, incubated for 30-60 minutes at 30°C to give <30% substrate turnover and terminated with 50 μ l yttrium silicate SPA beads (containing 3 mM of the respective unlabelled cyclic nucleotide for PDEs 9 and 11).
25 Plates are re-sealed and shaken for 20 minutes, after which the beads are allowed to settle for 30 minutes in the dark and then counted on a TopCount plate reader (Packard, Meriden, CT) Radioactivity units are converted to percent activity of an uninhibited control (100%), plotted against inhibitor concentration and inhibitor IC₅₀ values obtained using the 'Fit Curve' Microsoft Excel extension.

30 Effect of Specific PDE9 inhibitors on Insulin Resistance Syndrome in animals—Effects on Plasma Glucose, Triglyceride, Insulin, and cGMP Levels in ob/ob Mice.

Biological Data

Experimental Protocol

Test Compounds:

35 The PDE9 inhibitor compounds to be tested are solubilized in 10% DMSO/0.1% pluronics and dosed via oral gavage using mouse oral feeding needles (20 gauge, Popper & Sons,

Inc., New Hyde Park, NY). A volume of 4 ml/kg weight is administered for each dose. Compounds are tested at doses ranging from 1-50 mg/kg.

Experimental Animals:

Male *ob/ob* mice obtained from Jackson Laboratories (Bar Harbor, ME) are used in the studies at 6 to 10 weeks of age. Mice are housed five per cage and allowed free access to D11 mouse chow (Purina, Brentwood, MO) and water.

Experimental Protocol:

Mice are allowed to acclimate to the Pfizer animal facilities for one week prior to the start of the study. On day one, retro-orbital blood samples are obtained and plasma glucose is determined as described hereinafter. Mice are then sorted into groups of five such that mean plasma glucose concentrations for each group do not differ. On day one, mice are dosed with vehicle or a test PDE9 inhibitor compound only in the afternoon. Subsequently, mice are dosed twice a day on day 2-4 in the morning and in the afternoon. On day five, the mice receive an a.m. dose and are bled 3 hours later for plasma preparation for glucose and triglyceride analysis as described below. Terminal plasma samples are collected on day five following the retro-orbital sinus bleed as described below. Body weight is measured on days one and five of the study, and food consumption is assessed over the five day period.

Terminal Bleed and Tissue Collection:

On the morning of the last day of the study mice are dosed with test compound or vehicle at approximately 8:00 am. Three hours after dosing, 25 μ L of blood is obtained via the retro-orbital sinus and is added to 100 μ L of 0.025 percent heparinized-saline in Denville Scientific microtubes. The tubes are spun at the highest setting in a Beckman Microfuge 12 for 2 minutes. Plasma is collected for plasma glucose and triglyceride determination. The mice are then sacrificed by decapitation and about one milliliter of blood is collected in Becton-Dickinson Microtainer brand plasma separator tubes with lithium heparin. The tubes are spun in a Beckman Microfuge 12 at the maximum setting for five minutes. Plasma is collected in 1.5 ml Eppendorf tubes and snap frozen in liquid nitrogen. Plasma samples are stored at -80° C until analyzed.

Metabolite and Hormone Analysis:

Plasma glucose and triglycerides are measured using the Alcyon Clinical Chemistry Analyzer (Abbott Laboratories, Abbott Park, IL) using kits supplied by Abbott. Plasma cGMP is measured using the Biotrak enzyme-immunoassay system by Amersham (Piscataway, NJ). Via a similar technique the plasma insulin is assessed by the Mercodia ELISA Insulin kit by ALPCO (Uppsala, Sweden). All assays are conducted according to instructions provided by the manufacturers.

Results

Table 1 illustrates the changes in plasma glucose, triglyceride, and insulin levels over a five day period observed with Compound A, 5-(3-chloro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one.

Taken together, these experimental results in the hyperglycemic, insulin-resistant *ob/ob* mouse suggest that selective PDE9 inhibition improves metabolic parameters associated with IRS.

Table 1

Treatment	Plasma Glucose (mg/dl)	Plasma Triglyceride (mg/dl)	Plasma Insulin (pmol/ml)
Vehicle	370 ± 23	207 ± 9	12.0 ± 1.5
Compound A (10 mg/kg)	304 ± 17	155 ± 8	8.2 ± 1.5

The data in Table 1 are presented as mean ± standard error of the mean.


Table 2 illustrates the elevation of plasma cGMP produced by five day treatment with 5-(3-chloro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one.

Table 2

Treatment	Plasma cGMP (pmol/ml)
Vehicle	9.8 ± 0.5
Compound A (10 mg/kg)	16.8 ± 3.1

The data in Table 2 are presented as mean ± standard error of the mean.

General procedure for the Preparation of Examples 1 to 77



The carboxylic acids (80 μmol) were dissolved in a 3.75% solution of triethylamine in dimethylacetamide (400 μl) and administered to a 96 well plate. Carbonyldiimidazole (13 mg, 80 μmol) dissolved in pyridine (212 μl) was then added into each well, and the plates were left to stand at room temperature for 2 hours. A solution of 4-amino-5-isopropyl-2H-

32

pyrazole-3-carboxylic acid amide (13.5 mg, 80 μ mol) dissolved in dimethylacetamide (100 μ l) was then added, and the plates were sealed and heated to 70°C in an oven under nitrogen. This was maintained for 18 hours, upon which the plates were removed and allowed to cool to room temperature (2 hours). The solvent was removed using a
5 GENEVAC (45°C, 0.15mbar) over 5.5 hours. A solution of potassium t-butoxide (268 mg, 240 μ mol) in isopropylalcohol (0.5 ml) was added to each well, and the plates were sealed and transferred to an oven at 110°C under nitrogen. This was maintained for 15 hours, upon which the plates were removed and allowed to cool to room temperature (2 hours). The solvent was again removed using the GENEVAC (45°C, 0.15mbar) over 5.5 hours,
10 and a solution of p-toluenesulfonic acid (30 mg, 160 μ l) in isopropylalcohol (0.5 ml) was added to each well. The plates were left to stand at room temperature for 18 hours, and the solvent was removed using the GENEVAC (45°C, 0.15mbar) over 5.5 hours. The residues were dissolved in dimethylsulfoxide (450 μ l per well) and each compound was purified by preparative HPLC. The compounds were characterised by LC-MS analysis.

15

Preparative HPLC Conditions

Column : Phenomenex Luna C18, 5 μ m, 150 x 10mm id
Temperature : Ambient
20 Eluent A : 0.05% Diethylamine (aqueous)
Eluent B : Acetonitrile
Sample solvent : 90% dimethylsulfoxide in water
Initial pump conditions : A% 90, B% 10, flow 6 ml/minute
Detection : Gilston 119 uv detector – 225 nm
25 Injection volume – 600 μ l

30

Gradient Timetable

35

Time (min)	A%	B%	Flow (ml/min)
0.0	95	5	6
0.2	95	5	6
7.0	5	95	6
40 9.0	5	95	6
9.1	95	5	6
10.5	95	5	6

LC-MS ConditionsColumn : Phenomenex Luna C18, 5 μ m, 30 x 4.6 mm id.

Temperature : 40°C

5 Eluent A : 0.05% Diethylamine (aqueous)

Eluent B : Acetonitrile

Initial pump conditions : A% 90, B% 10, flow 3 ml/minute

Injection volume – 5 μ l10 Detection : Start range 210 nm, End range 280 nm, Range interval 5 nm, threshold
0.1 mAU, peakwidth 0.4 min.



Gradient Timetable

15	Time (min)	A%	B%	Flow (ml/min)	Pressure (bar)
	0.0	90	10	3	400
	2.2	5	95	3	400
	2.4	5	95	3	400
20	2.5	90	10	3	400






ELSD : Sedere Dedex 55, Temperature : 40°C, Gas Flow : 2.3bar






25 MS : Platform LC, ES+ Cone voltage : 26v, Capillary : 4.08kv
 ES- Cone voltage : -24v, Capillary : -3.58kv
 Blanket gas : 500 l/min, Temperature : 130°C

30







Example No.	Compound	Molecular Weight	Retention Time (min)
1	 5-(3-Chloro-benzyl)-3-isopropyl- 1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	302.1	1.95
2	 5-(3-Ethoxy-benzyl)-3-isopropyl- 1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	312.2	1.85

34






3	 5-Cyclohexylmethyl-3-isopropyl- 1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	274.2	1.92
4	 3-Isopropyl-5-(3-phenoxy-benzyl)- 1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	360.2	2.02
5	 3-Isopropyl-5-(2-trifluoromethyl- benzyl)-1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	302.1	1.85
6	 5-(3-Chloro-benzyl)-3-isopropyl- 1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	336.1	1.96
7	 5-(4-Chloro-benzyl)-3-isopropyl- 1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	302.1	1.91






8	 <p>5-(4-Benzyloxy-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	374.2	2.05
9	 <p>5-Biphenyl-4-ylmethyl-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	344.2	2.04
10	 <p>5-[2-(2-Chloro-phenyl)-ethyl]-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	316.1	1.93
11	 <p>3-Isopropyl-5-(2,4,6-trifluorobenzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	322.1	1.89
12	 <p>5-(3,5-Bis-trifluoromethyl-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	404.1	2.07

36






13	 3-Isopropyl-5-(3-trifluoromethoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	352.1	1.97
14	 5-(4-Butoxy-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	340.2	2.08
15	 3-Isopropyl-5-(3-methyl-butyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	248.2	1.80
16	 3-Isopropyl-5-(4-methylsulfanyl-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	314.2	1.83
17	 5-Ethyl-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	206.1	1.42
18	 3-Isopropyl-5-(3-methyl-butyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	234.2	1.63






37

	5-Isobutyl-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one		
19	 <p>3-Isopropyl-5-(4-methoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	298.1	1.77
20	 <p>5-(2,5-Dimethyl-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	296.2	1.93
21	 <p>5-Benzhydryl-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	344.2	2.07
22	 <p>3-Isopropyl-5-(9H-xanthen-9-yl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	358.1	2.08
23	 <p>5-(2-Fluoro-3-trifluoromethyl-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	354.1	1.95







24	 5-(2,4-Difluoro-benzyl)-3-isopropyl- 1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	304.1	1.82
25	 5-(2,3-Difluoro-benzyl)-3-isopropyl- 1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	304.1	1.86
26	 5-(4-Fluoro-benzyl)-3-isopropyl- 1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	286.1	1.80
27	 5-[2-(2-Imidazol-1-yl-ethoxy)- benzyl]-3-isopropyl-1,6-dihydro- pyrazolo[4,3-d]pyrimidin-7-one	378.2	1.44
28	 5-(5-Fluoro-2-trifluoromethyl- benzyl)-3-isopropyl-1,6-dihydro- pyrazolo[4,3-d]pyrimidin-7-one	354.1	1.98






39

29	 5-(2,6-Dichloro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	336.1	1.97
30	 5-(2-Chloro-6-methyl-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	332.1	2.03
31	 3-Isopropyl-5-(2-methoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	298.1	1.76
32	 5-Cyclopentyl-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	246.2	1.83
33	 5-(3,4-Difluoro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	304.1	1.83
34		339.2	1.73






	 <p>N-[1<i>R</i>-1-(3-Isopropyl-7-oxo-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2-phenyl-ethyl]-acetamide</p>		
35	 <p>5-(4-Methyl-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	282.2	1.83
36	 <p>3-Isopropyl-5-[2-(4-methoxy-phenyl)-1-phenyl-ethyl]-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	388.2	2.13
37	 <p>3-Isopropyl-5-napthalen-1-ylmethyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	318.2	2.05
38	 <p>5-Cyclopentylmethyl-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	260.2	1.77

41






39	 5-(2,6-Difluoro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	304.1	1.83
40	 5-(3-Methyl-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	282.2	1.84
41	 5-(2,4-Dimethyl-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	296.2	1.99
42	 5-(3-Fluoro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	286.1	1.82
43	 5-(2,3,6-Trifluoro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	322.1	1.91
44	 5-(4-chlorobenzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	318.1	1.97

	5-(4-Chloro-phenoxy-methyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one		
45	 3-Isopropyl-5-(4-phenoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	360.2	2.05
46	 5-(2-Chloro-6-fluoro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	320.1	1.88
47	 5-(2-Benzyloxy-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	374.2	2.04
48	 3-Isopropyl-5-(4-methyl-cyclohexylmethyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	288.2	2.02
49	 3-Isopropyl-5-(1 <i>R</i> -1-phenyl-propyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	296.2	2.09






43

50	 5-{2-[3-(4-Chloro-phenyl)- [1,2,4]oxadiazol-5-yl]-ethyl}-3- isopropyl-1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	384.1	2.05
51	 3-Isopropyl-5-pentyl-1,6-dihydro- pyrazolo[4,3-d]pyrimidin-7-one	248.2	1.83
52	 3-Isopropyl-5-(2-phenoxy-benzyl)- 1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	360.2	2.03
53	 5-(3,5-Dimethyl-benzyl)-3- isopropyl-1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	296.2	1.97
54	 5-(4-Cyclopentyloxy-3-methoxy- benzyl)-3-isopropyl-1,6-dihydro- pyrazolo[4,3-d]pyrimidin-7-one	382.2	1.95







44






55	 <p>3-Isopropyl-5-naphthalen-2-ylmethyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	318.2	1.97
56	 <p>5-(2,5-Dichlorophenylsulfanylmethyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	368.0	2.09
57	 <p>3-Isopropyl-5-(1S-1-phenyl-ethyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	282.2	1.95
58	 <p>3-Isopropyl-5-(2-methyl-butyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	248.2	1.78
59	 <p>5-(2,5-Difluoro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	304.1	1.86

45



60	 5-Benzyl-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	268.1	1.77
61	 3-Isopropyl-5-(4-methyl-pentyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	262.2	1.94
62	 5-(2-Cyclohexyl-ethyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	288.2	2.07
63	 5-(2-Chloro-4-fluoro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	320.1	1.92
64	 3-Isopropyl-5-[2-(4-trifluoromethyl-phenyl)-ethyl]-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	350.1	2.03

46


65	 <p>5-(2-Ethoxy-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	312.2	1.88
66	 <p>3-Isopropyl-5-phenoxy-methyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	284.1	1.85
67	 <p>5-(3-Methoxy-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	298.1	1.76
68	 <p>5-(3-Trifluoromethyl-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	336.1	1.99
69	 <p>5-(4-Isopropyl-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	310.2	2.04
70		304.1	1.89

	5-(3,5-Difluoro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one		
71	 5-(2,5-Dimethoxy-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	328.2	1.80
72	 5-(2,3-Dimethyl-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	312.2	2.04
73	 5-(3,4-Dichloro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	336.1	2.02
74	 5-(4-Trifluoromethyl-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	336.1	1.93
75	 5-(2-Methyl-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	282.2	1.85

48

76	 5-(2-Fluoro-benzyl)-3-isopropyl- 1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	286.1	1.80
77	 3-Isopropyl-5-(4-phenyl-butyl)-1,6- dihydro-pyrazolo[4,3-d]pyrimidin-7- one	310.2	1.98

49

General procedure for the Preparation of Examples 78 to 159

The carboxylic acids (80 μ mol) were dissolved in a 3.75% solution of triethylamine in dimethylacetamide (400 μ l) and administered to a 96 well plate. Carbonyldiimidazole (13 mg, 80 μ mol) dissolved in pyridine (212 μ l) was then added into each well, and the plates were left to stand at room temperature for 2 hours. A solution of 5-substituted-4-amino-pyrazole-3-carboxamide (80 μ mol) dissolved in dimethylacetamide (100 μ l) was then added, and the plates were sealed and heated to 70°C in an oven under nitrogen. This was maintained for 18 hours, upon which the plates were removed and allowed to cool to room temperature (2 hours). The solvent was removed using a GENEVAC (30°C, 0.15mbar) over 11 hours. A solution of potassium t-butoxide (268 mg, 240 μ mol) in isopropylalcohol (0.5 ml) was added to each well, and the plates were sealed and transferred to an oven at 110°C under nitrogen. This was maintained for 15 hours, upon which the plates were removed and allowed to cool to room temperature (2 hours). The solvent was again removed using the GENEVAC (30°C, 0.15mbar) over 11 hours, and a solution of p-toluenesulfonic acid (30 mg, 160 μ l) in isopropylalcohol (0.5 ml) was added to each well. The plates were left to stand at room temperature for 18 hours, and the solvent was removed using the GENEVAC (30°C, 0.15mbar) over 11 hours. The residues were dissolved in dimethylsulfoxide (450 μ l per well) and each compound was purified by preparative HPLC. The compounds were characterised by LC-MS analysis.

Preparative HPLC Conditions

Column : Phenomenex Luna C18, 5 μ m, 150 x 10 mm id
 Temperature : Ambient
 Eluent A : 0.05% Diethylamine (aqueous)
 Eluent B : Acetonitrile
 Sample solvent : 90% dimethylsulfoxide in water
 Initial pump conditions : A% 90, B% 10, flow 6 ml/minute
 Detection : Gilston 119 uv detector – 225 nm
 Injection volume – 600 μ l

Gradient Timetable

Time (min)	A%	B%	Flow (ml/min)
0.0	95	5	6
0.2	95	5	6
7.0	5	95	6
9.0	5	95	6
9.1	95	5	6
10.5	95	5	6

50

LC-MS ConditionsColumn : Phenomenex Luna C18, 5 μ m, 30 x 4.6 mm id.

Temperature : 40°C

5 Eluent A : 0.05% Diethylamine (aqueous)

Eluent B : Acetonitrile

Initial pump conditions : A% 90, B% 10, flow 3 ml/minute

Injection volume – 5 μ l

10 Detection : Start range 210 nm, End range 280 nm, Range interval 5 nm, threshold
0.1 mAU, peakwidth 0.4 min.

Gradient Timetable

	Time (min)	A%	B%	Flow (ml/min)	Pressure (bar)
15	0.0	90	10	3	400
	2.2	5	95	3	400
	2.4	5	95	3	400
	2.5	90	10	3	400

20



ELSD : Sedere Dedex 55, Temperature : 40°C, Gas Flow : 2.3bar

MS : Platform LC, ES+ Cone voltage : 26v, Capillary : 4.08kv






ES- Cone voltage : -24v, Capillary : -3.58kv

25






Blanket gas : 500 l/min, Temperature : 130°C

78	 <p>5-(2,6-Dichloro-benzyl)-3-pyridin-3-yl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	371.0	1.24
79	 <p>5-Cyclopropylmethyl-3-pyridin-3-yl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	267.1	0.92







51

80	 5-(2,6-Difluoro-benzyl)-3-pyridin-3-yl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	339.1	1.10
81	 5-(4-Methyl-cyclohexylmethyl)-3-pyridin-3-yl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	323.2	1.46
82	 5-(3-Chloro-benzyl)-3-pyridin-3-yl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	337.1	1.23
83	 5-(2-Ethoxy-benzyl)-3-pyridin-3-yl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	347.1	1.24
84	 5-(2-Phenoxy-benzyl)-3-pyridin-3-yl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	395.1	1.46






52

85	 <p>5-(2,3,5-Trifluoro-benzyl)-3-pyridin-3-yl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	357.1	1.18
86	 <p>5-(3-Fluoro-4-trifluoromethyl-benzyl)-3-pyridin-3-yl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	389.1	1.37
87	 <p>5-(5-Fluoro-2-trifluoromethyl-benzyl)-3-pyridin-3-yl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	389.1	1.29
88	 <p>5-(5-Bromo-2-methoxy-benzyl)-3-pyridin-3-yl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	411.0	1.30
89	 <p>5-(2-Benzyloxy-benzyl)-3-pyridin-3-yl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	409.2	1.42







53

90	 3-Butyl-5-(2-methyl-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	296.2	1.44
91	 3-Butyl-5-(2-methoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	312.2	1.36
92	 3-Butyl-5-(2-chloro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	316.1	1.45
93	 3-Butyl-5-(2-fluoro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	300.14	1.34
94	 3-Butyl-5-(2-chloro-6-fluoro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	334.1	1.47
95	 3-Butyl-5-(3,4-dichlorobenzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	350.1	1.56






54







	3-Butyl-5-(2,6-dichloro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one		
96	 <p>5-Butoxy-3-butyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	354.2	1.72
97	 <p>3-Butyl-5-cyclopropylmethyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	246.2	1.14
98	 <p>3-Butyl-5-(2,6-difluoro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	318.1	1.38
99	 <p>3-Butyl-5-(2-ethoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	326.2	1.48
100	 <p>3-Butyl-5-(2-benzyloxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	388.2	1.67






55






101	 <p>5-(2,4,5-Trifluoro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	322.1	1.36
102	 <p>5-(2,4-Dichloro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	336.1	1.54
103	 <p>5-(5-Bromo-2-methoxy-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	376.1	1.46
104	 <p>5-(2,3,6-Trichloro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	370.0	1.59
105	 <p>5-(3-Benzyloxy-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	374.2	1.59
106		220.1	0.96

56






	3-isopropyl-5-propyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one		
107	 <p>5-(2-Trifluoromethoxy-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	352.1	1.49
108	 <p>3-tert-Butyl-5-(3-chloro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	316.1	1.61
109	 <p>3-tert-Butyl-5-(5-bromo-2-methoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	390.1	1.68
110	 <p>3-Isobutyl-5-(2,4,5-trifluoro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	336.1	1.42
111	 <p>3-Isobutyl-5-(2-methyl-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	296.2	1.40

112	 <p>5-Cyclopentylmethyl-3-isobutyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	274.2	1.39
113	 <p>3,5-Diisobutyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	248.2	1.20
114	 <p>3-Isobutyl-5-(2-methoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	312.2	1.32
115	 <p>3-Isobutyl-5-(2-chloro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	316.1	1.41
116	 <p>3-Isobutyl-5-(2-fluoro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	300.1	1.31
117	 <p>3-Isobutyl-5-(2-chloro-3-fluorobenzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	334.1	1.44






	3-Isobutyl-5-(2-chloro-6-fluoro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one		
118	 3-Isopropyl-5-(2-methyl-butyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	262.2	1.35
119	 3-Isobutyl-5-(2-trifluoromethyl-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	350.1	1.53
120	 3-Isobutyl-5-(2,4-dichloro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	350.1	1.61
121	 3-Isobutyl-5-(2,6-dichloro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	350.1	1.54
122	 5-(4-Butoxy-benzyl)-3-Isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	354.2	1.69






123	 5-Cyclopropylmethyl-3-isobutyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	246.2	1.08
124	 3-Isobutyl-5-(2,6-difluoro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	318.1	1.34
125	 3-Isobutyl-5-(3-chloro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	316.1	1.44
126	 3-Isobutyl-5-(2-ethoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	326.2	1.44
127	 3-Isobutyl-5-(2-phenoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	374.2	1.65






60






128	 <p>3-Isobutyl-5-(2,3,5-trifluoro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	336.1	1.43
129	 <p>3-Isobutyl-5-(5-bromo-2-methoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	390.1	1.53
130	 <p>3-Isobutyl-5-(2-benzyloxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	388.2	1.64
131	 <p>5-[2-(2-Imidazol-1-yl-ethoxy)-benzyl]-3-isobutyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	392.2	1.12
132	 <p>3-Isobutyl-5-(2,3,6-trichloro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	384.0	1.66

61






133	 3-Isobutyl-5-(3-benzyloxy-benzyl)- 1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	388.2	1.64
134	 5-(2,3-Dihydro-benzofuran-5- ylmethyl)-3-isobutyl-1,6-dihydro- pyrazolo[4,3-d]pyrimidin-7-one	324.2	1.25
135	 3-Cyclopentyl-5-(2,4,5-trifluoro- benzyl)-1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	348.1	1.51
136	 3-Cyclopentyl-5-(2-methyl-benzyl)- 1,6-dihydro-pyrazolo[4,3- d]pyrimidin-7-one	308.2	1.52
137	 3-Cyclopentyl-5-isobutyl-1,6- dihydro-pyrazolo[4,3-d]pyrimidin-7- one	260.2	1.33

138	 3-Cyclopentyl-5-(2-methoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	324.2	1.44
139	 3-Cyclopentyl-5-(2-chloro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	328.1	1.52
140	 3-Cyclopentyl-5-(2-fluoro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	312.1	1.42
141	 3-Cyclopentyl-5-(2-chloro-6-fluoro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	346.1	1.53
142	 3-Cyclopentyl-5-(2-methyl-butyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	274.2	1.49



143	 3-Cyclopentyl-5-(2-trifluoromethyl-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	362.1	1.62
144	 3-Cyclopentyl-5-(2,4-dichloro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	362.1	1.70
145	 3-Cyclopentyl-5-(2,6-dichloro-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	362.1	1.61
146	 5-(4-Butoxy-benzyl)-3-cyclopentyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	366.2	1.80
147	 3-Cyclopentyl-5-cyclopropylmethyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	258.2	1.22


148	 3-Cyclopentyl-5-(2,6-difluorobenzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	330.1	1.44
149	 3,5-Dicyclopentyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	272.2	1.52
150	 3-Cyclopentyl-5-(3-chlorobenzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	328.1	1.55
151	 3-Cyclopentyl-5-(2,5-dimethoxybenzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	354.2	1.41
152	 3-Cyclopentyl-5-(2-ethoxybenzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one	338.2	1.55

65

153	 <p>3-Cyclopentyl-5-(2-phenoxybenzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	386.2	1.75
154	 <p>3-Cyclopentyl-5-(3-fluoro-4-(trifluoromethyl)benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	380.1	1.67
155	 <p>3-Cyclopentyl-5-(2-benzyloxybenzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	400.2	1.73
156	 <p>3-Cyclopentyl-5-[2-(2-imidazo-1-ylethoxy)benzyl]-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	404.2	1.20
157	 <p>3-Cyclopentyl-5-(3-benzyloxybenzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	400.2	1.73

66

158	 <p>3-Cyclopentyl-5-propyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	246.2	1.19
159	 <p>3-Cyclopentyl-5-(2,3-dihydro-benzofuran-5-ylmethyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one</p>	336.2	1.35

Example 160**3-Cyclopentyl-5-(2-trifluoromethoxy-benxyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one.**

5

5-Cyclopentyl-4-[2-(2-trifluoromethoxy-phenyl)-acetylamino]-1H-pyrazole-3-carboxylic acid amide (120 mg, 0.303 mmol) and potassium tert-butoxide (102 mg, 0.909 mmol) were suspended in isopropylalcohol (5 ml) and the reaction was heated to reflux, under nitrogen, for 18 hours. The reaction mixture was concentrated under reduced pressure and the residue was partitioned between ethyl acetate (20 ml) and water (20 ml). The aqueous phase was removed, acidified to pH 2 with 2N HCl, and extracted with ethyl acetate (2 x 15 ml). The combined organic extracts were washed with saturated sodium carbonate solution (3 x 10 ml), dried over MgSO₄, concentrated under reduced pressure and the residue was purified by flash column chromatography on silica gel eluting with dichloromethane : methanol (95 : 5, by volume) to give 3-cyclopentyl-5-(2-trifluoromethoxy-benxyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one (21 mg) as an off-


67

white solid. ^1H NMR (400MHz, DMSO- d_6): δ = 7.36-7.41 (2H, m), 7.29-7.36 (2H, m), 3.97-4.03 (2H, brs), 2.39-2.45 (1H, m, partially masked by solvent), 1.82-1.94 (2H, m), 1.66-1.79 (2H, m), 1.58-1.65 (2H, m), 1.49-1.58 (2H, m) ppm. LRMS (electrospray): m/z $[\text{M}-\text{H}]^+$ 377.

5

Example 161

3-isobutyl-5-(2-trifluoromethoxy-benxyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one.




5-isobutyl-4-[2-(2-trifluoromethoxy-phenyl)-acetylamino]-1H-pyrazole-3-carboxylic acid amide (140 mg, 0.365 mmol) and potassium tert-butoxide (123 mg, 1.09 mmol) were suspended in isopropylalcohol (6 ml) and the reaction was heated to reflux, under nitrogen, for 18 hours. The reaction mixture was concentrated under reduced pressure and the residue was partitioned between ethyl acetate (20 ml) and water (20 ml). The aqueous phase was removed, acidified to pH 2 with 2N HCl, and extracted with ethyl acetate (2 x 15 ml). The combined organic extracts were washed with saturated sodium carbonate solution (3 x 10 ml), dried over MgSO_4 , concentrated under reduced pressure and the residue was purified by flash column chromatography on silica gel eluting with dichloromethane : methanol (95 : 5, by volume) to give 3-isobutyl-5-(2-trifluoromethoxy-benxyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one (27 mg) as an off-white solid. ^1H NMR (400MHz, CDCl_3): δ = 8.64-8.74 (1H, brs), 7.22-7.41 (4H, m, partially masked by solvent), 4.15 (2H, s), 2.79-2.84 (2H, d), 2.13-2.23 (1H, m), 0.92-1.00 (6H, d) ppm. LRMS (electrospray): m/z $[\text{M}+\text{H}]^+$ 367, $[\text{M}-\text{H}]^+$ 365.

Example 162

3-Pyridin-3-yl-5-(2-trifluoromethoxy-benxyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one.


68



5-Pyridin-3-yl-4-[2-(2-trifluoromethoxy-phenyl)-acetylamino]-1H-pyrazole-3-carboxylic acid amide (345 mg, 0.85 mmol) and potassium tert-butoxide (286 mg, 2.55 mmol) were suspended in isopropylalcohol (5 ml) and the reaction was heated to 55°C under nitrogen for 18 hours. The reaction mixture was concentrated under reduced pressure and the residue was partitioned between ethyl acetate (20 ml) and water (20 ml). The aqueous phase was removed, acidified to pH 2 with 2N HCl, and extracted with ethyl acetate (2 x 15 ml) and dichloromethane (2 x 15 ml). The combined organic extracts were dried over MgSO₄, concentrated under reduced pressure and the residue was purified by flash column chromatography on silica gel eluting with a solvent gradient of dichloromethane : methanol (99 : 1 changing to 95 : 5, by volume). The product was triturated with methanol (3 ml), dichloromethane (3 ml) and diethylether (3 ml) to give 3-pyridin-3-yl-5-(2-trifluoromethoxy-benxyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one (13mg) as an off-white solid. ¹H NMR (400MHz, CD₃OD): δ = 9.34 (1H, brs), 8.57-8.61 (1H, d), 8.43-8.48 (1H, m), 7.32-7.47 (5H, m), 4.18 (2H, s) ppm. LRMS (electrospray): m/z [M-H]⁺ 386.

Preparation 1

4-Methyl-3-oxo-pentanoic acid ethyl ester.



Sodium pellets (3.39 g, 148 mmol) were dissolved in ethanol (100 ml) under nitrogen at room temperature and a solution of diethyloxalate (20 ml, 147 mmol) in 3-methyl-2-butanone (18.9 ml, 177 mmol) was added dropwise at room temperature over 30 minutes. The reaction was diluted with ethanol (100 ml), heated to 60°C and stirred at this temperature for 2 hours. After cooling to room temperature the reaction was poured onto ice-cold 2N HCl (200 ml) and extracted with diethylether (300 ml) and ethyl acetate (300 ml). The combined organic extracts were dried over MgSO₄, concentrated under reduced pressure and the residue was purified by flash column chromatography on silica gel eluting with a solvent gradient of pentane : ethyl acetate (99 : 1 changing to 95 : 5, by

69

volume) to give 4-methyl-3-oxo-pentanoic acid ethyl ester (23.8 g) as a yellow oil. ^1H NMR (400MHz, CDCl_3): δ = 14.40-14.80 (1H, brs), 6.40 (1H, s), 4.30-4.39 (2H, quart), 2.60-2.71 (1H, quin), 1.35-1.40 (3H, t), 1.15-1.20 (6H, d) ppm. LRMS (electrospray): m/z $[\text{M}-\text{H}]^+$ 185.

5

Preparation 2

5-Isopropyl-1H-pyrazol-3-carboxylic acid ethyl ester.



Hydrazine hydrate (6.6 ml, 134 mmol) was added to a solution of 4-methyl-3-oxo-pentanoic acid ethyl ester (23.8 g, 188 mmol) in ethanol (100 ml) at room temperature under nitrogen. The reaction was allowed to proceed at room temperature for 18 hours, and the solvent was removed under reduced pressure. The residue was partitioned between dichloromethane (300 ml) and water (300 ml) and the aqueous phase was removed. The organic phase was washed with water (2 x 200 ml), dried over MgSO_4 and concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel eluting with a solvent gradient of pentane : ethyl acetate (4 : 1 changing to 2 : 1, by volume) to give 5-isopropyl-1H-pyrazol-3-carboxylic acid ethyl ester (18.9 g) as a white solid. ^1H NMR (400MHz, CDCl_3): δ = 10.80-10.95 (1H, brs), 6.61 (1H, s), 4.33-4.40 (2H, quart), 2.98-3.08 (1H, quin), 1.35-1.41 (3H, t), 1.24-1.32 (6H, d) ppm. LRMS (electrospray): m/z $[\text{M}-\text{H}]^+$ 181.

20

Preparation 3

5-Isopropyl-1H-pyrazol-3-carboxylic acid.



5-Isopropyl-1H-pyrazol-3-carboxylic acid ethyl ester (18.9 g, 104 mmol) and 1M NaOH solution (260 ml, 259 mmol) were dissolved in 1,4-dioxan (300 ml), the reaction was heated to 50°C under nitrogen and stirred for 3 hours. The reaction mixture was cooled, adjusted to pH 2 using concentrated hydrochloric acid and the solvent was removed under reduced pressure. The residual solid was azeotroped with toluene (2 x 30 ml), dissolved in ethyl acetate (500 ml) and washed with water (200 ml). The aqueous phase was removed, extracted with ethyl acetate (2 x 200 ml) and the combined organic extracts were dried over MgSO_4 . The solvent was removed under reduced pressure and the

30


70

residue was azeotroped with dichloromethane (2 x 50 ml) to give 5-isopropyl-1H-pyrazol-3-carboxylic acid (14.7 g) as a white solid. ^1H NMR (400MHz, DMSO- D_6): δ = 12.50-13.30 (2H, brs), 6.42 (1H, s), 2.84-2.94 (1H, quin), 1.15-1.19 (6H, d) ppm. LRMS (electrospray): m/z $[\text{M}-\text{H}]^+$ 153.

5

Preparation 4

5-Isopropyl-4-nitro-1H-pyrazol-3-carboxylic acid.




5-Isopropyl-1H-pyrazol-3-carboxylic acid (5 g, 32.5 mmol) was added portionwise to concentrated sulfuric acid (25 ml) at room temperature with stirring. The reaction mixture was then heated to 60°C and concentrated nitric acid (70%, 6ml, 90 mmol) was added dropwise, keeping the temperature at 60°C. The reaction was then stirred at 60°C for 3 hours, cooled to room temperature and poured onto 50 ml of ice with stirring. After 15 minutes the white precipitate was isolated by filtration, washed with water and dried under reduced pressure to give 5-isopropyl-4-nitro-1H-pyrazol-3-carboxylic acid (5.2 g) as a white solid. ^1H NMR (400MHz, DMSO- D_6): δ = 13.86-13.93 (1H, brs), 13.50-13.80 (1H, brs), 3.39-3.52 (1H, m), 1.18-1.30 (6H, d) ppm. LRMS (electrospray): m/z $[\text{M}-\text{H}]^+$ 198.

15

Preparation 5

5-Isopropyl-4-nitro-1H-pyrazol-3-carboxylic acid amide.



Oxalyl chloride (6.8 ml, 77.6 mmol) was added dropwise to a suspension of 5-isopropyl-4-nitro-1H-pyrazol-3-carboxylic acid (5.15 g, 25.9 mmol) in dichloromethane (80 ml) containing dimethylformamide (0.1 ml) under nitrogen at 0°C. The reaction was stirred at 0°C for 1 hours, allowed to warm to room temperature and stirred for a further 2 hours. The solvent was removed under reduced pressure, the residue was dissolved in toluene (100 ml) and ammonia gas was bubbled into the solution for 2 hours. The reaction was stirred under nitrogen at room temperature for 18 hours, concentrated under reduced pressure and the residue was dissolved in hot methanol (300 ml). The resultant precipitate was filtered and the filtrate was concentrated under reduced pressure. The residue was azeotroped with water (300 ml), concentrated to approximately 80 ml under reduced pressure and the precipitate was isolated by filtration. This was washed with


25

30


71

water and dried under to give 5-isopropyl-4-nitro-1H-pyrazol-3-carboxylic acid amide (3.1g) as an orange solid. ^1H NMR (400MHz, DMSO- D_6): δ = 7.94-7.99 (1H, brs), 7.68-7.72 (1H, brs), 3.45-3.55 (1H, m), 1.24-1.30 (6H, d) ppm. LRMS (electrospray): m/z $[\text{M}+\text{Na}]^+$ 221, $[\text{M}-\text{H}]^+$ 197.

5

Preparation 64-Amino-5-isopropyl-1H-pyrazol-3-carboxylic acid amide.

5-Isopropyl-4-nitro-1H-pyrazol-3-carboxylic acid amide (3 g, 15.1 mmol) and 10% palladium on carbon (500 mg) in ethanol (30 ml) were stirred under hydrogen (50 psi) at room temperature for 18 hours. The reaction mixture was filtered and the solid was washed with methanol (50 ml), dichloromethane (50 ml), ethanol (50 ml) and ethyl acetate (50 ml). The filtrate was concentrated under reduced pressure and the residue was purified by flash column chromatography on silica gel eluting with dichloromethane : methanol (9 : 1, by volume) to give 4-amino-5-isopropyl-1H-pyrazol-3-carboxylic acid amide (2.6 g) as an off-white solid. ^1H NMR (400MHz, DMSO- D_6): δ = 12.20-12.30 (1H, brs), 7.02-7.14 (1H, brs), 6.85-6.95 (1H, brs), 4.30-4.46 (2H, brs), 2.90-3.00 (1H, m), 1.15-1.21 (6H, d) ppm. LRMS (electrospray): m/z $[\text{M}-\text{H}]^+$ 167, $[\text{2M}-\text{H}]^+$ 335. Anal. Found C, 49.86; H, 7.21; N, 33.07. $\text{C}_7\text{H}_{12}\text{N}_4\text{O}$ requires C, 49.99; H, 7.19; N, 33.31%.


Preparation 73-Oxo-heptanoic acid ethyl ester.

Sodium pellets (3.82 g, 166 mmol) were dissolved in ethanol (100 ml) under nitrogen at room temperature and a solution of diethyloxalate (22.6 ml, 166 mmol) hexan-2-one (20 g, 198 mmol) was added dropwise at room temperature over 30 minutes. The reaction was diluted with ethanol (100 ml), heated to 60°C and stirred at this temperature for 2 hours. After cooling to room temperature the reaction was poured onto ice-cold 2N HCl (200 ml) and extracted with diethylether (300 ml) and ethyl acetate (300 ml). The combined organic extracts were dried over MgSO_4 , concentrated under reduced pressure and the residue was purified by flash column chromatography on silica gel eluting with a solvent gradient of pentane : ethyl acetate (99 : 1 changing to 95 : 5, by volume) to give 3-oxo-heptanoic acid ethyl ester (30.3 g) as an orange oil. ^1H NMR (400MHz, CDCl_3): δ =

14.30-14.80 (1H, brs), 6.37 (1H, s), 4.30-4.39 (2H, quart), 2.43-2.50 (2H, t), 1.59-1.62 (2H, quin), 1.31-1.40 (5H, t + m), 0.86-0.97 (3H, t) ppm. LRMS (electrospray): m/z $[M-H]^+$ 199.

Preparation 8


5 5-Butyl-1H-pyrazol-3-carboxylic acid ethyl ester.



Hydrazine hydrate (7.75 ml, 157 mmol) was added dropwise to a solution of 3-oxoheptanoic acid ethyl ester (30 g, 150 mmol) in ethanol (100 ml) at room temperature under nitrogen. The reaction was heated to 50°C and allowed to proceed at this temperature for 18 hours, and the solvent was removed under reduced pressure. The residue was partitioned between dichloromethane (300 ml) and water (300 ml) and the aqueous phase was removed. The organic phase was washed with water (2 x 200 ml), dried over $MgSO_4$ and concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel eluting with pentane : ethyl acetate (8 : 1, by volume) to give 5-butyl-1H-pyrazol-3-carboxylic acid ethyl ester (24 g) as a yellow oil. 1H NMR (400MHz, $CDCl_3$): δ = 10.58-10.78 (1H, brs), 6.62 (1H, s), 4.35-4.40 (2H, quart), 2.63-2.70 (2H, t), 1.60-1.67 (2H, quin), 1.35-1.42 (5H, t + m), 0.90-0.96 (3H, t) ppm. LRMS (electrospray): m/z $[M-H]^+$ 195.

Preparation 9

20 5-Butyl-1H-pyrazol-3-carboxylic acid.




5-Butyl-1H-pyrazol-3-carboxylic acid ethyl ester (24 g, 122 mmol) and 1M NaOH solution (305 ml, 306 mmol) were dissolved in 1,4-dioxan (300 ml), the reaction was heated to 55°C under nitrogen and stirred for 2 hours. The reaction mixture was cooled, adjusted to pH 2 using concentrated hydrochloric acid and the solvent was removed under reduced pressure. The residual solid was dissolved in ethyl acetate (300 ml) and washed with water (300 ml). The aqueous phase was removed, extracted with ethyl acetate (300 ml) and the combined organic extracts were dried over $MgSO_4$. The solvent was removed

under reduced pressure and the residue was azeotroped with dichloromethane (2 x 50 ml) to give 5-butyl-1H-pyrazol-3-carboxylic acid (22.6 g) as a white solid. ¹H NMR (400MHz, DMSO-D₆): δ = 12.50-13.00 (2H, brs), 6.41 (1H, s), 2.47-2.57 (2H, t), 1.46-1.56 (2H, quin), 1.19-1.29 (2H, sext), 1.15-1.19 (3H, t) ppm. LRMS (electrospray): m/z [M-H]⁺ 167. Anal. Found C, 57.01; H, 7.23; N, 16.50. C₈H₁₂N₂O₂ requires C, 57.13; H, 7.19; N, 16.66%.

Preparation 10

5-Butyl-4-nitro-1H-pyrazol-3-carboxylic acid.




- 10 5-Butyl-1H-pyrazol-3-carboxylic acid (22.6 g, 134 mmol) was added portionwise to concentrated sulfuric acid (100 ml) at room temperature with stirring. The reaction mixture was then heated to 60°C and concentrated nitric acid (70%, 23.7 ml, 376 mmol) was added dropwise, keeping the temperature at 60°C. The reaction was then stirred at 60°C for 3 hours, cooled to room temperature and poured onto 50 ml of ice with stirring.
- 15 After 15 minutes the pale yellow precipitate was isolated by filtration, washed with water and dried under reduced pressure to give 5-butyl-4-nitro-1H-pyrazol-3-carboxylic acid (21.9 g) as a pale yellow solid. ¹H NMR (400MHz, DMSO-D₆): δ = 2.83-2.92 (2H, t), 1.56-1.64 (2H, quin), 1.22-1.36 (2H, sext), 0.84-0.90 (3H, t) ppm. LRMS (electrospray): m/z [M-H]⁺ 212. Anal. Found C, 30.19; H, 5.41; N, 13.12. C₈H₁₁N₃O₄. 6mol H₂O requires C, 29.91; H, 7.22; N, 13.08%.
- 20

Preparation 11

5-Butyl-4-nitro-1H-pyrazol-3-carboxylic acid amide.


74



Oxalyl chloride (12.3 ml, 141 mmol) was added dropwise to a suspension of 5-butyl-4-nitro-1H-pyrazol-3-carboxylic acid (10 g, 46.9 mmol) in dichloromethane (100 ml) containing dimethylformamide (0.5 ml) under nitrogen at 0°C. The reaction was stirred at 0°C for 1 hours, allowed to warm to room temperature and stirred for a further 2 hours. The solvent was removed under reduced pressure, the residue was dissolved in toluene (100 ml) and ammonia gas was bubbled into the solution for 2 hours. The reaction was stirred under nitrogen at room temperature for 18 hours, concentrated under reduced pressure and the residue was purified by flash column chromatography on silica gel eluting with dichloromethane : methanol (9 : 1, by volume) to give 5-butyl-4-nitro-1H-pyrazol-3-carboxylic acid amide (3.1 g) as an orange solid. ¹H NMR (400MHz, DMSO-D₆): δ = 7.87-7.96 (1H, brs), 7.57-7.66 (1H, brs), 2.83-2.90 (2H, t), 1.56-1.63 (2H, quin), 1.24-1.36 (2H, sext), 0.84-0.92 (3H, t) ppm. LRMS (electrospray): m/z [M-H]⁺ 211. Anal. Found C, 44.66; H, 5.56; N, 25.50. C₈H₁₂N₄O₃. 0.23mol H₂O requires C, 44.41; H, 5.80; N, 25.90%.

Preparation 12


4-Amino-5-butyl-1H-pyrazol-3-carboxylic acid amide.



5-butyl-4-nitro-1H-pyrazol-3-carboxylic acid amide (3.1 g, 17.0 mmol) and 10% palladium on carbon (600 mg) in ethanol (50 ml) were stirred under hydrogen (50psi) at room temperature for 18 hours. The reaction mixture was filtered and the solid was washed with methanol (50 ml), dichloromethane (50 ml), ethanol (50 ml) and ethyl acetate (50 ml). The filtrate was concentrated under reduced pressure and the residue was purified by flash column chromatography on silica gel eluting with a solvent gradient of dichloromethane : methanol (95 : 5 changing to 90 : 10, by volume) to give 4-amino-5-butyl-1H-pyrazol-3-carboxylic acid amide (2.37g) as an an orange solid. ¹H NMR (400MHz, DMSO-D₆): δ = 12.24-12.32 (1H, brs), 7.02-7.14 (1H, brs), 6.80-6.95 (1H, brs),

75

4.28-4.46 (2H, brs), 2.39-2.50 (2H, t, partially masked by solvent), 1.45-1.56 (2H, quin), 1.22-1.35 (2H, sext), 0.83-0.90 ppm. LRMS (electrospray): m/z $[M-H]^+$ 181, $[2M-H]^+$ 363. Anal. Found C, 52.58; H, 7.80; N, 30.56. $C_8H_{14}N_4O$ requires C, 52.73; H, 7.74; N, 30.75%.

Preparation 135 4,4-Dimethyl-3-oxo-pentanoic acid ethyl ester.

Sodium pellets (4.6 g, 200 mmol) were dissolved in ethanol (165 ml) under nitrogen at room temperature and a solution of diethyloxalate (27.2 ml, 200 mmol) in tert-butyl-methyl ketone (20.1 g, 200 mmol) was added dropwise at room temperature over 15 minutes.

10 The reaction was diluted with ethanol (100 ml), heated to 60°C and stirred at this temperature for 2 hours. After cooling to room temperature the reaction was stirred for 64 hours, poured onto ice-cold 2N HCl (200 ml) and extracted with diethylether (3 x 200 ml). The combined organic extracts were washed with water, dried over $MgSO_4$ and concentrated under reduced pressure to give 4,4-dimethyl-3-oxo-pentanoic acid ethyl ester (36.7g) as a yellow oil. 1H NMR (400MHz, $CDCl_3$): δ = 6.48 (1H, s), 4.26-4.37 (2H, quart), 1.29-1.38 (3H, t), 1.17 (9H, s) ppm. LRMS (electrospray): m/z $[M+Na]^+$ 223, $[M-H]^+$ 199.

Preparation 145-tert-Butyl-1H-pyrazol-3-carboxylic acid ethyl ester.

20 Hydrazine hydrate (9.5 ml, 180 mmol) was added to a solution of 4,4-dimethyl-3-oxo-pentanoic acid ethyl ester (36.7 g, 180 mmol) in ethanol (188 ml) at room temperature under nitrogen. The reaction was allowed to proceed at room temperature for 2 hours, and the solvent was removed under reduced pressure. The residue was partitioned

25 between dichloromethane (500 ml) and water (400 ml) and the aqueous phase was removed. The organic phase was washed with brine (200 ml), dried over $MgSO_4$ and concentrated under reduced pressure to give 5-tert-butyl-1H-pyrazol-3-carboxylic acid ethyl ester (30.6 g) as a white solid. 1H NMR (400MHz, $CDCl_3$): δ = 14.50-14.90 (1H, brs), 6.45 (1H, s), 4.25-4.31 (2H, quart), 1.27-1.36 (3H, t), 1.16 (9H, s) ppm. LRMS (thermospray): m/z $[M+H]^+$ 197. Anal. Found C, 61.12; H, 8.20; N, 14.28. $C_{10}H_{16}N_2O_2$ requires C, 61.20; H, 8.22; N, 14.27%.

30

76

Preparation 155-tert-Butyl-1H-pyrazol-3-carboxylic acid.

5-tert-Butyl-1H-pyrazol-3-carboxylic acid ethyl ester (20 g, 100 mmol) and 1M NaOH solution (250 ml, 250 mmol) were dissolved in 1,4-dioxan (300 ml), the reaction was heated to 60°C under nitrogen and stirred for 2.5 hours. The reaction was then cooled to room temperature and stirred for a further 18 hours. The reaction mixture was adjusted to pH 2 using concentrated hydrochloric acid, extracted with ethyl acetate (4 x 200 ml) and the combined organic extracts washed with brine (100 ml). The organic phase was dried over MgSO₄ and the solvent was removed under reduced pressure to give 5-tert-butyl-1H-pyrazol-3-carboxylic acid (14.7 g) as a pale yellow solid. ¹H NMR (400MHz, DMSO-D₆): δ = 12.80-12.88 (2H, brs), 6.41 (1H, s), 1.11 (9H, s) ppm. LRMS (electrospray): m/z [M+H]⁺ 169, [M+Na]⁺ 191, [M-H]⁺ 167.

Preparation 165-tert-Butyl-4-nitro-1H-pyrazol-3-carboxylic acid.

5-tert-Butyl-1H-pyrazol-3-carboxylic acid (5 g, 29.7 mmol) was added portionwise to concentrated sulfuric acid (25 ml) at room temperature with stirring. The reaction mixture was then heated to 60°C and concentrated nitric acid (70%, 5.15 ml) was added dropwise, keeping the temperature at 60°C. The reaction was then stirred at 60°C for 2.5 hours, cooled to room temperature and poured onto 50 ml of ice with stirring. After 15 minutes the white precipitate was isolated by filtration, washed with water and dried under reduced pressure to give 5-tert-butyl-4-nitro-1H-pyrazol-3-carboxylic acid (6.0g) as a white solid. ¹H NMR (400MHz, DMSO-D₆): δ = 13.50-13.88 (2H, brs), 1.13 (9H, s) ppm. LRMS (electrospray): m/z [M+Na]⁺ 236, [M-H]⁺ 212.

Preparation 175-tert-Butyl-4-nitro-1H-pyrazol-3-carboxylic acid amide.

77



Oxalyl chloride (10.2 ml, 117 mmol) was added dropwise to a suspension of 5-tert-butyl-4-nitro-1H-pyrazol-3-carboxylic acid (6 g, 28 mmol) in dichloromethane (55 ml) containing dimethylformamide (0.1 ml) under nitrogen at 0°C. The reaction was stirred at 0°C for 0.5
 5 hours, allowed to warm to room temperature and stirred for a further 1.5 hours. The solvent was removed under reduced pressure, the residue was azeotroped with dichloromethane (50 ml) and the residue was dissolved in dichloromethane (100 ml). Ammonia gas was bubbled into the solution for 45 minutes and the reaction was stirred under nitrogen at room temperature for 18 hours, concentrated under reduced pressure
 10 and the residue was dissolved in ethyl acetate (250 ml). After washing with water (100 ml) and brine (100 ml) the organic phase was filtered, the filtrate was dried over MgSO₄ and concentrated under reduced pressure to give 5-tert-butyl-4-nitro-1H-pyrazol-3-carboxylic acid amide (4.0 g) as a light brown solid. ¹H NMR (400MHz, DMSO-D₆): δ = 12.90-13.08 (1H, brs), 7.78-7.86 (1H, brs), 7.49-7.60 (1H, brs), 1.30 (9H, s) ppm. LRMS
 15 (electrospray): m/z [M+H]⁺ 213, [M+Na]⁺ 235, [M-H]⁺ 211.

Preparation 18

4-Amino-5-tert-butyl-1H-pyrazol-3-carboxylic acid amide.



5-tert-Butyl-4-nitro-1H-pyrazol-3-carboxylic acid amide (4.6 g, 21 mmol) and 10%
 20 palladium on carbon (300 mg) in ethanol (80 ml) was stirred under hydrogen (60psi) at room temperature for 18 hours. The reaction mixture was filtered and the filtrate was concentrated under reduced pressure. The residue was pre-absorbed onto silica gel and purified by flash column chromatography eluting with a solvent gradient of dichloromethane : methanol (100 : 0 changing to 95 : 5 then 90 : 10, by volume) to give 4-
 25 amino-5-tert-butyl-1H-pyrazol-3-carboxylic acid amide (2.96 g) as an off-white solid, which was a mixture of rotamers. ¹H NMR (400MHz, DMSO-D₆): δ = 12.10-12.20 (0.75H, brs), 11.75-11.85 (0.25H, brs), 7.04-7.16 (1.5H, brs), 6.88-6.96 (0.5H, brs), 4.27-4.59 (2H, 2xbrs), 1.12 (9H, s) ppm. LRMS (electrospray) : m/z [M+H]⁺ 183, [M+Na]⁺ 205, [M-H]⁺ 181. Anal. Found C, 52.45; H, 7.84; N, 30.62. C₈H₁₄N₄O requires C, 52.73; H, 7.74; N,
 30 30.75%.

78

Preparation 195-Methyl-3-oxo-hexanoic acid ethyl ester.

Sodium pellets (4.6 g, 200 mmol) were dissolved in ethanol (165 ml) under nitrogen at room temperature and a solution of diethyloxalate (13.5 ml, 100 mmol) in isobutylmethyl ketone (30 ml, 200 mmol) was added dropwise at room temperature over 20 minutes. The reaction was heated to 60°C and stirred at this temperature for one hour. After cooling to room temperature the reaction was poured onto ice-cold 2N HCl (200 ml) and extracted with diethylether (4 x 200 ml). The combined organic extracts were washed with water, dried over MgSO₄ and concentrated under reduced pressure to give 5-methyl-3-oxo-hexanoic acid ethyl ester (20 g) as a yellow oil. ¹H NMR (400MHz, CDCl₃): δ = 14.40-14.70 (1H, brs), 6.27 (1H, s), 4.25-4.32 (2H, quart), 2.26-2.31 (2H, d), 2.02-2.18 (1H, m), 1.29-1.34 (3H, t), 0.89-0.94 (6H, d) ppm. LRMS (thermospray): m/z [M+NH₄]⁺ 218.

15

Preparation 205-Isobutyl-1H-pyrazol-3-carboxylic acid ethyl ester.


Hydrazine hydrate (5.7 ml, 115 mmol) was added to a solution of 5-methyl-3-oxo-hexanoic acid ethyl ester (22 g, 110 mmol) in ethanol (113 ml) at room temperature under nitrogen. The reaction was allowed to proceed at room temperature for 18 hours, and the solvent was removed under reduced pressure. The residue was partitioned between dichloromethane (400 ml) and water (400 ml) and the aqueous phase was removed. The organic phase was washed with brine (200 ml), water (200 ml), dried over MgSO₄ and concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel eluting with a solvent gradient of pentane : ethyl acetate (1 : 0 changing to 6 : 1, 5 : 1, 4 : 1, 3 : 1, 2 : 1 and finally 1 : 1, by volume) to give 5-isobutyl-1H-pyrazol-3-carboxylic acid ethyl ester (16.5 g) as a white solid. ¹H NMR (400MHz, CDCl₃): δ = 11.60-12.60 (1H, brs), 6.53 (1H, s), 4.26-4.35 (2H, quart), 2.48-2.54 (2H, d), 1.80-1.90 (1H, m), 1.25-1.31 (3H, t), 0.81-0.88 (6H, d) ppm. LRMS (thermospray): m/z [M+H]⁺ 197, [2M+H]⁺ 393. Anal. Found C, 61.49; H, 8.30; N, 14.24. C₁₀H₁₈N₂O₂ requires C, 61.20; H, 8.22; N, 14.27%.

79

Preparation 215-Isobutyl-1H-pyrazol-3-carboxylic acid.

- 5-Isobutyl-1H-pyrazol-3-carboxylic acid ethyl ester (16.2 g, 83 mmol) and 1M NaOH solution (173 ml, 173 mmol) were dissolved in 1,4-dioxan (260 ml) and the reaction was stirred at room temperature under nitrogen for 64 hours. The reaction mixture was adjusted to pH 7 using concentrated hydrochloric acid, and concentrated under reduced pressure. Water (500 ml) was added, the slurry was adjusted to pH 1 with concentrated hydrochloric acid and the aqueous phase was extracted with ethyl acetate (5 x 300 ml). The combined organic extracts were dried over MgSO_4 and the solvent was removed under reduced pressure to give 5-isobutyl-1H-pyrazol-3-carboxylic acid (10 g) as a white solid. ^1H NMR (400MHz, DMSO-D_6): δ = 12.72-12.90 (1H, brs), 6.39 (1H, s), 2.39-2.43 (2H, d), 1.77-1.86 (1H, m), 0.78-0.83 (6H, d) ppm. LRMS (electrospray): m/z $[\text{M}+\text{Na}]^+$ 191, $[\text{2M}+\text{Na}]^+$ 359, $[\text{M}-\text{H}]^+$ 167, $[\text{2M}-\text{H}]^+$ 335.

15


Preparation 225-Isobutyl-4-nitro-1H-pyrazol-3-carboxylic acid.

- 5-Isobutyl-1H-pyrazol-3-carboxylic acid (5 g, 29.7 mmol) was added portionwise to concentrated sulfuric acid (25 ml) at room temperature with stirring. The reaction mixture was then heated to 60°C and concentrated nitric acid (70%, 5.15 ml) was added dropwise, keeping the temperature at 60°C. The reaction was then stirred at 60°C for 3 hours, cooled to room temperature and poured onto 50 ml of ice with stirring. The resultant white precipitate was isolated by filtration, washed with water and dried under reduced pressure to give 5-isobutyl-4-nitro-1H-pyrazol-3-carboxylic acid (6.4 g) as a white solid. ^1H NMR (400MHz, DMSO-D_6): δ = 2.71-2.76 (2H, d), 1.88-2.00 (1H, m), 0.80-0.87 (6H, d) ppm. LRMS (thermospray): m/z $[\text{M}+\text{NH}_4]^+$ 231, $[\text{M}-\text{H}]^+$ 212. LRMS (electrospray): m/z $[\text{M}-\text{H}]^+$ 212, $[\text{2M}-\text{H}]^+$ 425. Anal. Found C, 42.54; H, 5.18; N, 18.63. $\text{C}_8\text{H}_{11}\text{N}_3\text{O}_4$. 0.7mol H_2O requires C, 42.55; H, 5.54; N, 18.61%.

Preparation 23

- 30 5-Isobutyl-4-nitro-1H-pyrazol-3-carboxylic acid amide.

80



Oxalyl chloride (10 ml, 115 mmol) was added dropwise to a suspension of 5-isobutyl-4-nitro-1H-pyrazol-3-carboxylic acid (5.6 g, 26 mmol) in dichloromethane (70 ml) containing dimethylformamide (0.1 ml) under nitrogen at 0°C. The reaction was stirred at 0°C for 0.5
 5 hours, allowed to warm to room temperature and stirred for a further 2 hours. The solvent was removed under reduced pressure, the residue was azeotroped with dichloromethane (3 x 50 ml) and the residue was dissolved in toluene (100 ml). Ammonia gas was bubbled into the solution for 2 hours and the reaction was stirred under nitrogen at room temperature for 18 hours, concentrated under reduced pressure and the residue was
 10 suspended in methanol (250 ml). After filtration, the filtrates were concentrated under reduced pressure, the residue was dissolved in ethyl acetate (400 ml) and washed with water (50 ml). The organic phase was filtered, the filtrate was dried over MgSO₄ and concentrated under reduced pressure. The filtered solid and residue from the filtrates were combined to give 5-isobutyl-4-nitro-1H-pyrazol-3-carboxylic acid amide (4.8 g) as an
 15 off-white solid. ¹H NMR (400MHz, DMSO-D₆): δ = 13.61-13.81 (1H, brs), 7.80-7.96 (1H, brs), 7.50-7.66 (1H, brs), 2.70-2.76 (2H, d), 1.90-2.01 (1H, m), 0.83-0.88 (6H, d) ppm. LRMS (electrospray): m/z [M+Na]⁺ 235, [2M+Na]⁺ 447, [M-H]⁺ 211, [2M-H]⁺ 423. Anal. Found C, 45.12; H, 5.68; N, 26.31. C₈H₁₂N₄O₃ requires C, 45.28; H, 5.70; N, 26.40%.

Preparation 24

20 4-Amino-5-isobutyl-1H-pyrazol-3-carboxylic acid amide.




5-Isobutyl-4-nitro-1H-pyrazol-3-carboxylic acid amide (4.7 g, 22 mmol) and 10% palladium on carbon (300 mg) in ethanol (80 ml) was stirred under hydrogen (60psi) at room temperature for 4 hours, and held under nitrogen for 64 hours. The reaction mixture was
 25 filtered and the filtrate was concentrated under reduced pressure of dichloromethane : methanol (100 : 0 changing to 95 : 5 then 90 : 10, by volume) to give 4-amino-5-isobutyl-1H-pyrazol-3-carboxylic acid amide (3.8 g) as an off-white solid, which was a mixture of rotamers. ¹H NMR (400MHz, DMSO-D₆): δ = 12.20-12.28 (1H, brs), 7.00-7.10 (1.34H, brs), 6.80-6.85 (0.66H, brs), 4.27-4.40 (2H, brs), 2.27-2.36 (2H, d), 1.78-1.88 (1H, m),
 30 0.77-0.84 (6H, d) ppm. LRMS (electrospray) : m/z [M+H]⁺ 183, [M+Na]⁺ 205. Anal. Found

81

C, 52.27; H, 7.78; N, 30.59. $C_8H_{14}N_4O$ requires C, 52.73; H, 7.76; N, 30.75%.

Preparation 25


1-Cyclopentylethanone.



- 5 Concentrated sulfuric acid (22.4 ml, 420 mmol) was slowly added to a solution of chromium trioxide (26.3 g, 263 mmol) dissolved in water (50 ml) at room temperature. After 10min this solution was added to 1-cyclopentylethanol (20 g, 175 mmol) dissolved in acetone (450 ml) maintaining the temperature below 35°C. The addition was continued until a bright orange colour persisted for 10 minutes. The reaction mixture was quenched
- 10 with isopropyl alcohol to destroy excess chromic acid and it was then neutralised to pH 5 with the portionwise addition of sodium bicarbonate. After filtration the filtrate was concentrated under reduced pressure (to 50 ml) and extracted with diethylether (3 x 300 ml). The combined organic extracts were dried over $MgSO_4$ and concentrated under reduced pressure to give 1-cyclopentylethanone (16.7 g) as a colourless oil. 1H NMR
- 15 (400MHz, $CDCl_3$): δ 2.80-2.90 (1H, quin), 2.18 (3H, s), 1.53-1.86 (8H, 2xm) ppm.

Preparation 26

3-Cyclopentyl-3-oxo-propionic acid ethyl ester.



- Sodium pellets (3.1 g, 135 mmol) were dissolved in ethanol (100 ml) under nitrogen at
- 20 room temperature and a solution of diethyloxalate (18.4 ml, 135mmol) and 1-cyclopentylethanone (16.7 g, 149 mmol) was added dropwise at room temperature over 30 minutes. The reaction was diluted with ethanol (100 ml), heated to 60°C and stirred at this temperature for 2 hours. After cooling to room temperature the reaction was poured onto ice-cold 2N HCl (200 ml) and extracted with diethylether (300 ml) and ethyl acetate
- 25 (300 ml). The combined organic extracts were dried over $MgSO_4$, concentrated under reduced pressure and the residue was purified by flash column chromatography on silica gel eluting with pentane : ethyl acetate (6 : 1, by volume) to give 3-cyclopentyl-3-oxo-propionic acid ethyl ester (23.8 g) as an orange oil. 1H NMR (400MHz, $CDCl_3$): δ = 14.38-14.65 (1H, brs), 6.83 (1H, s), 4.30-4.39 (2H, quart), 2.82-2.92 (1H, quin), 1.83-1.96 (2H,
- 30 m), 1.57-1.83 (6H, 2xm), 1.33-1.40 (3H, t) ppm. LRMS (electrospray): m/z $[M-H]^+$ 211.

Preparation 27

5-Cyclopentyl-1H-pyrazol-3-carboxylic acid ethyl ester.


82



Hydrazine hydrate (5.8 ml, 117 mmol) was added to a solution of 3-cyclopentyl-3-oxopropionic acid ethyl ester (23.7 g, 112 mmol) in ethanol (100 ml) at room temperature under nitrogen. The reaction was allowed to proceed at room temperature for 18 hours, then heated to 50°C and held at this temperature for 4 hours. The solvent was removed under reduced pressure and the residue was partitioned between dichloromethane (300 ml) and water (300 ml) and the aqueous phase was removed. The organic phase was washed with water (2 x 200 ml), dried over MgSO₄ and concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel eluting with a solvent gradient of pentane : ethyl acetate (4 : 1, by volume) to give 5-cyclopentyl-1H-pyrazol-3-carboxylic acid ethyl ester (17.1 g) as a pale yellow solid. ¹H NMR (400MHz, CDCl₃): δ = 10.40-10.60 (1H, brs), 6.58 (1H, s), 4.30-4.38 (2H, quart), 3.01-3.10 (1H, quin), 2.00-2.10 (2H, m), 1.56-1.80 (6H, 2xm), 1.33-1.39 (3H, t) ppm. LRMS (electrospray): m/z [M+H]⁺ 209, [M+Na]⁺ 231. Anal. Found C, 63.40; H, 7.75; N, 13.41. C₁₁H₁₆N₂O₂ requires C, 63.44; H, 7.74; N, 13.45%.

Preparation 28

5-Cyclopentyl-1H-pyrazol-3-carboxylic acid.



5-Cyclopentyl-1H-pyrazol-3-carboxylic acid ethyl ester (17.1 g, 82 mmol) and 1M NaOH solution (205 ml, 205 mmol) were dissolved in 1,4-dioxan (300 ml) and the reaction was heated to 50°C under nitrogen and stirred for 3 hours. The reaction mixture was cooled, adjusted to pH 2 using concentrated hydrochloric acid and the solvent was removed under reduced pressure. The residual solid was azeotrope with toluene (2 x 30 ml), dissolved in ethyl acetate (500 ml) and washed with water (200 ml). The aqueous phase was removed, extracted with ethyl acetate (2 x 200 ml) and the combined organic extracts were dried over MgSO₄. The solvent was removed under reduced pressure and the residue was azeotrope with dichloromethane (2 x 50 ml) to give 5-cyclopentyl-1H-pyrazol-3-carboxylic acid (13 g) as a white solid. ¹H NMR (400MHz, DMSO-D₆): δ = 12.75-12.88 (2H, brs), 6.43 (1H, s), 2.97-3.08 (1H, quin), 1.91-2.02 (2H, m), 1.50-1.76 (6H, 2xm) ppm.

83

LRMS (electrospray): m/z $[M-H]^+$ 179. Anal. Found C, 59.72; H, 6.74; N, 15.37. $C_9H_{12}N_2O_2$ requires C, 59.99; H, 6.71; N, 15.55%.

Preparation 295-Cyclopentyl-4-nitro-1H-pyrazol-3-carboxylic acid.

5-Cyclopentyl-1H-pyrazol-3-carboxylic acid (13 g, 72.1 mmol) was added portionwise to concentrated sulfuric acid (75 ml) at room temperature with stirring. The reaction mixture was then heated to 60°C and concentrated nitric acid (70%, 12.7 ml, 202 mmol) was added dropwise, keeping the temperature at 60°C. The reaction was then stirred at 60°C for 3 hours, cooled to room temperature and poured onto 50 ml of ice with stirring. After 15 minutes the precipitate was isolated by filtration, washed with water and dried under reduced pressure to give 5-cyclopentyl-4-nitro-1H-pyrazol-3-carboxylic acid (7.1 g) as a yellow solid. 1H NMR (400MHz, DMSO- D_6): δ = 14.00-14.41 (1H, brs), 13.28-13.85 (1H, brs), 3.20-3.56 (1H, brs, partially masked by solvent), 1.96-2.10 (2H, m), 1.54-1.80 (6H, 2xm) ppm. LRMS (electrospray): m/z $[M-H]^+$ 224, $[2M-H]^+$ 449. Anal. Found C, 43.83; H, 5.35; N, 16.94. $C_9H_{11}N_3O_4 \cdot 1.2mol H_2O$ requires C, 43.80; H, 5.47; N, 17.02%.

Preparation 305-Cyclopentyl-4-nitro-1H-pyrazol-3-carboxylic acid amide.

Oxalyl chloride (7.65 ml, 87.7 mmol) was added dropwise to a suspension of 5-isopropyl-4-nitro-1H-pyrazol-3-carboxylic acid (6.58 g, 29.2 mmol) in dichloromethane (100 ml) containing dimethylformamide (0.5 ml) under nitrogen at 0°C. The reaction was stirred at 0°C for 1 hours, allowed to warm to room temperature and stirred for a further 2 hours. The solvent was removed under reduced pressure, the residue was azeotroped with dichloromethane (2 x 50 ml) and dissolved in toluene (100 ml). Ammonia gas was bubbled into the solution for 2 hours and the reaction was stirred under nitrogen at room temperature for 18 hours, concentrated under reduced pressure and purified by flash column chromatography on silica gel eluting with a solvent gradient of dichloromethane : methanol (95 : 5 changing to 90 : 10, by volume) to give 5-cyclopentyl-4-nitro-1H-pyrazol-

84

3-carboxylic acid amide (5.48 g) as a yellow solid. ^1H NMR (400MHz, DMSO- D_6): δ = 13.67-13.79 (1H, brs), 7.88-8.03 (1H, brs), 7.59-7.77 (1H, brs), 3.46-3.60 (1H, quin), 1.97-2.11 (2H, m), 1.58-1.81 (6H, 2xm) ppm. LRMS (electrospray): m/z $[\text{M}-\text{H}]^+$ 223, $[2\text{M}-\text{H}]^+$ 447. Anal. Found C, 56.12; H, 7.39; N, 27.55. $\text{C}_9\text{H}_{12}\text{N}_4\text{O}_3$. 0.2mol acetone requires C, 56.01; H, 7.44; N, 27.22%.

Preparation 31

4-Amino-5-cyclopentyl-1H-pyrazol-3-carboxylic acid amide.




5-cyclopentyl-4-nitro-1H-pyrazol-3-carboxylic acid amide (4.48 g, 20 mmol) and 10% palladium on carbon (800 mg) in ethanol (50 ml) were stirred under hydrogen (50psi) at room temperature for 18 hours. The reaction mixture was filtered through arboceel and the solid was washed with ethanol (50 ml), methanol (50 ml), dichloromethane (50 ml), and ethyl acetate (50 ml). The filtrate was concentrated under reduced pressure and the residue was purified by flash column chromatography on silica gel eluting with dichloromethane : methanol (9 : 1, by volume) to give 4-amino-5-cyclopentyl-1H-pyrazol-3-carboxylic acid amide (4.0 g) as an off-white solid which was a mixture of rotamers. ^1H NMR (400MHz, DMSO- D_6): δ = 12.20-12.31 (0.75H, brs), 11.78-11.87 (0.25H, brs), 7.02-7.18 (1.5H, brs), 6.80-6.93 (0.5H, brs), 4.22-4.56 (2H, 2xbrs), 2.92-3.02 (1H, quin), 1.79-1.96 (2H, m), 1.48-1.78 (6H, 2xm) ppm. LRMS (electrospray): m/z $[\text{M}-\text{H}]^+$ 193. Anal. Found C, 56.12; H, 7.39; N, 27.55. $\text{C}_9\text{H}_{14}\text{N}_4\text{O}$. 0.2mol acetone requires C, 56.01; H, 7.44; N, 27.22%.

25

Preparation 32

(3-Benzyloxy-phenyl)-acetic acid benzyl ester.


85



3-Hydroxy-phenyl-acetic acid (15.3 g, 101 mmol), benzyl bromide (36.2 g, 202 mmol) and potassium carbonate (29.2 g, 202 mmol) were suspended in dimethylformamide (300 ml) and the reaction was heated to reflux under nitrogen for 44 hours. The reaction mixture was cooled, filtered and the filtrate was concentrated under reduced pressure. The residue was partitioned between ethyl acetate (200 ml) and water (200 ml), and the aqueous phase was extracted with ethyl acetate (2 x 200 ml). The combined organic extracts were washed with brine (200 ml), dried over Na₂SO₄ and the solvent was removed under reduced pressure. The residue was purified by flash column chromatography on silica gel eluting with pentane : ethyl acetate (95 : 5, by volume) to give (3-benzyloxy-phenyl)-acetic acid benzyl ester (10.7 g) as a white solid..

Preparation 33

(3-Benzyloxy-phenyl)-acetic acid.




1N Sodium hydroxide solution (35 ml, 35 mmol) was added to a solution of (3-benzyloxy-phenyl)-acetic acid benzyl ester (5.3 g, 16 mmol) in methanol (350 ml) at room temperature under nitrogen. The reaction was heated to reflux for 2 hours, and the solvent was removed under reduced pressure. The residue was dissolved in water (500 ml) and extracted with ether (3 x 350 ml). The aqueous phase was acidified to pH 1 with concentrated hydrochloric acid and the resultant precipitate was isolated by filtration and dried under vacuum to give (3-benzyloxy-phenyl)-acetic acid (3.08 g) as a white solid. mp 127-129°C. ¹H NMR (400MHz, CDCl₃): δ = 7.26-7.43 (5H, m), 7.20-7.26 (1H, m, partially masked by solvent), 6.84-6.96 (3H, m + s), 5.04 (2H, s), 3.62 (2H, s) ppm. LRMS (electrospray): m/z [M-H]⁺ 241. Anal. Found C, 74.21; H, 5.82. C₁₅H₁₄O requires C, 74.36; H, 5.82%.

86

Preparation 34(4-hydroxy-3-methoxy-phenyl)-acetic acid methyl ester.


Concentrated sulfuric acid (12 ml) was added to a solution of (4-hydroxy-3-methoxy-phenyl)-acetic acid (22.5 g, 123 mmol) in methanol (450 ml) at room temperature, and the reaction was heated to 90°C for 2.45 hours. The reaction was then cooled to room temperature and stirred for 18 hours, and the solvent was removed under reduced pressure. The residue was suspended in ice water (300 ml) and extracted with diethylether (2 x 300 ml). The combined organic extracts were washed with saturated sodium bicarbonate solution (2 x 100 ml), brine (100 ml), dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel eluting with a solvent gradient of cyclohexane : ethyl acetate (80 : 20 changing to 70 : 30, 60 : 40 and finally 1 : 1, by volume) to give (4-hydroxy-3-methoxy-phenyl)-acetic acid methyl ester (23 g) as a yellow oil. ¹H NMR (400MHz, CDCl₃): δ = 6.82-6.85 (1H, d), 6.80 (1H, s), 6.76-6.79 (1H, d), 5.49 (1H, s), 3.86 (3H, s), 3.66 (3H, s), 3.53 (2H, s) ppm. LRMS (electrospray): m/z [M+Na]⁺ 219.

Preparation 35(4-Cyclopentyloxy-3-methoxy-phenyl)-acetic acid methyl ester.

Cyclopentanol (7.7 ml, 85 mmol) and triphenylphosphine (28 g, 107 mmol) were added to a solution of (4-hydroxy-3-methoxy-phenyl)-acetic acid methyl ester (14 g, 71 mmol) in tetrahydrofuran (280 ml) under nitrogen at 0°C. Diethylazodicarboxylate (15.7 ml, 100 mmol) was then added dropwise and the reaction was allowed to warm to room temperature and stirred for 44 hours. The solvent was removed under reduced pressure, pentane (200 ml) was added and the suspension was filtered. The filtrate was concentrated under reduced pressure and purified by flash column chromatography on

87

silica gel eluting with a solvent gradient of cyclohexane : ethyl acetate (90 : 10 changing to 85 : 15, by volume) to give (4-cyclopentyloxy-3-methoxy-phenyl)-acetic acid methyl ester (12.4 g) as a colourless oil. ¹H NMR (400MHz, CD₃OD): δ = 6.79-6.85 (2H, m), 6.73-6.79 (1H, d), 4.73-4.79 (1H, brs), 3.79 (3H, s), 3.64 (3H, s), 3.53 (2H, s), 1.74-1.89 (6H, m), 1.56-1.67 (2H, m) ppm. LRMS (electrospray): m/z [M+Na]⁺ 287. Anal. Found C, 68.01; H, 7.74. C₁₅H₂₀O₄ requires C, 68.16; H, 7.63%.


Preparation 36(4-Cyclopentyloxy-3-methoxy-phenyl)-acetic acid.

- 10 Sodium hydroxide (4.75 g, 119 mmol) was added to a solution of (4-cyclopentyloxy-3-methoxy-phenyl)-acetic acid methyl ester (12.4 g, 46.9 mmol) in methanol (100 ml) / water (100 ml) and the reaction was stirred at room temperature for 3.5 hours. The methanol was removed under reduced pressure and the aqueous phase was washed with diethylether (100 ml) then acidified to pH 2 using concentrated hydrochloric acid. This
- 15 was then extracted with ethyl acetate (2 x 200 ml) and the combined organic extracts were washed with brine (100 ml), dried over Na₂SO₄ and concentrated under reduced pressure to give (4-cyclopentyloxy-3-methoxy-phenyl)-acetic acid (11.1 g) as a white solid. ¹H NMR (400MHz, CD₃OD): δ = 6.87 (1H, s), 6.81-6.86 (1H, d), 6.76-6.80 (1H, d), 4.75-4.79 (1H, brs), 3.78 (3H, s), 3.49 (2H, s), 1.71-1.89 (6H, m), 1.56-1.64 (2H, m) ppm.
- 20 LRMS (electrospray): m/z [M-H]⁺ 249, [2M-H]⁺ 499. Anal. Found C, 67.15; H, 7.25. C₁₄H₁₈O₄ requires C, 67.18; H, 7.25%.

25

Preparation 372,4-Dimethyl-phenyl-acetic acid.


88



2,4-Dimethylbenzylcyanide (70 g, 0.48 mol) was mixed with water (134 ml) and concentrated sulfuric acid (106 ml, 1.98 mol) was added slowly. The reaction was heated to reflux for 3 hours, then cooled to room temperature over 18 hours. The mixture was
5 poured onto crushed ice (500ml), stirred for one hour and the resulting precipitate was isolated by filtration. After washing with water the solid was dissolved in 1.2M sodium hydroxide solution (500 ml), extracted with dichloromethane (2 x 250 ml) and the aqueous phase was treated with decolourising carbon (2 g) at reflux for 10 min and filtered hot through hyflo supercel. The filtrate was then acidified with concentrated hydrochloric acid
10 and the resulting precipitate was isolated by filtration, washed with water and dried under vacuum to give 2,4-dimethyl-phenyl-acetic acid (52.6 g) as a white solid. ¹H NMR (250MHz, CD₃OD/D₂O): δ = 6.88-7.03 (3H, m), 3.48-3.68 (2H, s), 2.23 (6H, s) ppm.

Preparation 38

Benzene sulfonic acid 2-chloro-ethyl ester.




15

2-Chloroethanol (1168 g, 975 mol) and benzene sulfonyl chloride (2780 g, 2015 mol) were stirred together at -5°C and pyridine (2158 g, 2200 mol) was added over a 3 hours period, maintaining the temperature below 0°C. The reaction was stirred for a further 3 hours at -5°C to 0°C and was then allowed to warm to room temperature over 18 hours.
20 After pouring into a mixture of ice (10 liters) and water (10 liters) the reaction was stirred for 15 minutes, extracted with ether (10 liters) and the organic phase was washed with 5N HCl (2 x 2 liters) and water (2 x 4 liters). It was then dried over MgSO₄ and concentrated under reduced pressure to give benzene sulfonic acid 2-chloro-ethyl ester (1921 g) as an orange oil. ¹H NMR (250MHz, CDCl₃): δ = 7.78-8.02 (2H, m), 7.58-7.78 (3H, m), 4.20-4.45
25 (2H, t), 3.60--3.81 (2H, t) ppm.

Preparation 39


2-Hydroxy-phenyl-acetic acid ethyl ester.



2-Hydroxy-phenyl-acetic acid (30.4 g, 0.2 mol) was dissolved in chloroform (200 ml) and thionyl chloride (50 ml, 0.2 mol) was added. The reaction was gently refluxed for 2 hours, upon which the mixture was concentrated under reduced pressure. The residue was slowly poured into ethanol (200 ml) maintaining a temperature of 10°C to 20°C. The solvent was removed under reduced pressure and the residue was purified by thermal distillation to give 2-hydroxy-phenyl-acetic acid ethyl ester (31.6 g) as a yellow oil. Bp 146-150°C. ν_{\max} (thin film) 1710 cm^{-1} (C=O, ester).

Preparation 40

10 [2-(2-Chloro-ethoxy)-phenyl]-acetic acid ethyl ester.




50% Sodium hydride in mineral oil (8.11 g, 169 mmol) was added portionwise to a solution of 2-hydroxy-phenyl-acetic acid ethyl ester (30.4 g, 169 mmol) in dimethylformamide (100 ml). After the initial effervescence had ended the reaction was heated to 100°C for 10 minutes and was cooled to room temperature. A solution of benzene sulfonic acid 2-chloro-ethyl ester (37.2 g, 169 mmol) in dimethylformamide (5 ml) was then added and the reaction was heated to 100°C for one hour, and allowed to cool to room temperature over 18 hours. The reaction mixture was partitioned between diethylether (300 ml) and water (300 ml) and the organic phase was removed and washed with water (100 ml), dried over MgSO_4 and the solvent was removed under reduced pressure. The residue was purified by thermal distillation to give [2-(2-chloro-ethoxy)-phenyl]-acetic acid ethyl ester (22.0 g) as a pale yellow oil. Bp 170°C to 180°C. ν_{\max} (thin film) 1735 cm^{-1} (C=O, ester); no O-H stretch. Anal. Found C, 59.35; H, 6.29. $\text{C}_{12}\text{H}_{15}\text{ClO}_3$ requires C, 59.38; H, 6.23%.

25

Preparation 41

[2-(2-Imidazol-1-yl-ethoxy)-phenyl]-acetic acid.


90



[2-(2-Imidazol-1-yl-ethoxy)-phenyl]-acetic acid ethyl ester (3.5 g, 113 mmol) was stirred in 50% aqueous hydrochloric acid (20 ml) at 100°C for 6 hours. After cooling to room temperature the solvent was removed under reduced pressure and the residue was recrystallised from isopropylalcohol to give [2-(2-Imidazol-1-yl-ethoxy)-phenyl]-acetic acid (2.73 g) as a white solid. Mp 146-147°C. ν_{\max} (thin film) 3410 (O-H), 1722 cm^{-1} (C=O, acid). Anal. Found C, 54.89; H, 5.25; N, 9.80. $\text{C}_{13}\text{H}_{14}\text{N}_2\text{O}_3$. 1 mol HCl requires C, 55.22; H, 5.35; N, 9.91%.

Preparation 42

10 5-Cyclopentyl-4-[2-(2-trifluoromethoxy-phenyl)-acetylamino]-1H-pyrazole-3-carboxylic acid amide.




Carbonyldiimidazole (84 mg, 0.515 mmol) was added to a solution of 2-trifluoromethoxy-phenyl-acetic acid (113 mg, 0.515 mmol) in tetrahydrofuran (4 ml) under nitrogen at room temperature, and the mixture was stirred for 3 hours. 4-Amino-5-cyclopentyl-1H-pyrazole-3-carboxylic acid amide (100 mg, 0.515 mmol) was then added and the reaction was stirred for 18 hours. The reaction mixture was diluted with water (20 ml), acidified to pH 2 with 2N HCl and extracted with ethyl acetate (2 x 20 ml). The combined organic extracts were dried over MgSO_4 and concentrated under reduced pressure to give 5-cyclopentyl-4-[2-(2-trifluoromethoxy-phenyl)-acetylamino]-1H-pyrazole-3-carboxylic acid amide (120 mg) as an off-white solid. LRMS (electrospray): m/z $[\text{M}+\text{H}]^+$ 397, $[\text{M}-\text{H}]^+$ 395.

25

Preparation 43

5-Isobutyl-4-[2-(2-trifluoromethoxy-phenyl)-acetylamino]-1H-pyrazole-3-carboxylic acid amide.


91



Carbonyldiimidazole (84 mg, 0.515 mmol) was added to a solution of 2-trifluoromethoxy-phenyl-acetic acid (113 mg, 0.515 mmol) in tetrahydrofuran (4 ml) under nitrogen at room temperature, and the mixture was stirred for 3 hours. 4-Amino-5-isobutyl-1H-pyrazol-3-carboxylic acid amide (100 mg, 0.515 mmol) was then added and the reaction was stirred for 18 hours. The reaction mixture was diluted with water (20 ml), acidified to pH2 with 2N HCl and extracted with ethyl acetate (2 x 20 ml). The combined organic extracts were dried over MgSO₄ and concentrated under reduced pressure to give 5-isobutyl-4-[2-(2-trifluoromethoxy-phenyl)-acetylamino]-1H-pyrazole-3-carboxylic acid amide (142 mg) as an off-white solid. LRMS (electrospray): m/z [M+H]⁺ 385, [M-H]⁺ 383.

Preparation 44

5-Pyridine-3-yl-4-[2-(2-trifluoromethoxy-phenyl)-acetylamino]-1H-pyrazole-3-carboxylic acid amide.



15


Carbonyldiimidazole (144 mg, 0.886 mmol) was added to a solution of 2-trifluoromethoxy-phenyl-acetic acid (195 mg, 0.886 mmol) in tetrahydrofuran (5 ml) under nitrogen at room temperature, and the mixture was stirred for one hour. 4-Amino-5-cyclopropyl-1H-pyrazol-3-carboxylic acid amide (180 mg, 0.886 mmol) was then added and the reaction was stirred for 18 hours. The reaction mixture was diluted with brine (20 ml) and extracted with ethyl acetate (2 x 20 ml). The combined organic extracts were dried over MgSO₄ and concentrated under reduced pressure to give 5-pyridine-3-yl-4-[2-(2-trifluoromethoxy-phenyl)-acetylamino]-1H-pyrazole-3-carboxylic acid amide (345 mg) as an off-white solid. LRMS (electrospray): m/z [M+Na]⁺ 428, [M-H]⁺ 404.

25

CLAIMS

1. A method of treating insulin resistance syndrome in a mammal comprising
 5 administering to said mammal a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said PDE9 inhibitor, prodrug, solvate or salt.

2. The method of claim 1 wherein said cGMP PDE9 inhibitor is a compound of the formula (I)



10 or a pharmaceutically acceptable salt, solvate or prodrug thereof, wherein:

R^1 is H or (C_1-C_6) alkyl;

R^2 is (C_1-C_6) alkyl, straight chain or branched chain, (C_3-C_7) cycloalkyl or heteroaryl;

R^3 is (C_1-C_6) alkyl, straight chain or branched chain, optionally substituted by 1-2 groups

15 each independently selected from Ar, (C_3-C_7) cycloalkyl, OAr, SAr, $NC(O)(C_1-C_6)$ alkyl, heteroaryl, xanthene, and naphthalene;

Ar is a group of formula



wherein R^4 , R^5 and R^6 are each independently selected from H, halo, phenoxy, phenyl,

20 CF_3 , OCF_3 , $S(C_1-C_6)$ alkyl, (C_1-C_6) alkyl, $O(C_1-C_6)$ alkyl, said alkyl optionally substituted by a heteroaryl group or by a phenyl group, wherein said phenyl group is optionally substituted by 1-3 groups selected from halo, CF_3 , OCF_3 and (C_1-C_6) alkyl; or wherein R^4 and R^5 may combine to form a (C_2-C_3) alkyl link, wherein said link may optionally incorporate a heteroatom selected from O, S and N; and

25 heteroaryl is aromatic 5-6 membered heterocycle containing 1-3 heteroatoms, each independently selected from O, S and N, said heterocycle optionally substituted by 1-3 substituents, each independently selected from (C_1-C_6) alkyl, halo and phenyl, said phenyl optionally substituted by 1-3 groups selected from halo and (C_1-C_6) alkyl;

with the proviso that when R^1 is $-CH_3$, R^2 cannot be $-CH_2CH_2CH_3$.


30 3. The method of claim 2 wherein R^1 is H or CH_3 ; R^2 is (C_3-C_4) alkyl,

cyclopentyl or pyridinyl; R^3 is (C_1-C_3) alkyl, optionally substituted by 1-2 groups selected from Ar, (C_3-C_7) cycloalkyl and heteroaryl; R^4 , R^5 and R^6 are each independently selected from H, halo, phenoxy, phenyl, CF_3 , OCF_3 , $S(C_1-C_6)$ alkyl, (C_1-C_6) alkyl, $O(C_1-C_6)$ alkyl; said alkyl in the definition of R^4 , R^5 and R^6 is optionally substituted by a heteroaryl group or by
 5 a phenyl group optionally substituted by 1-3 groups selected from halo, CF_3 , OCF_3 and (C_1-C_6) alkyl; or wherein R^4 and R^5 may combine to form a C_2 alkyl link, said link incorporating an O atom; and heteroaryl is an aromatic 5-6 membered heterocycle containing at least 2 nitrogen atoms, said heterocycle optionally substituted by 1-3
 10 substituents, each independently selected from (C_1-C_6) alkyl, halo and phenyl, said phenyl in the definition of heterocycle optionally substituted by 1-3 groups selected from halo and (C_1-C_6) alkyl.

4. The method of claim 1 comprising administering to said mammal 5-(3-chloro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one, a prodrug thereof or a pharmaceutically acceptable salt thereof or of said prodrug.

15 5. A method of treating type 2 diabetes in a mammal comprising administering to said mammal a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said PDE9 inhibitor, prodrug, solvate or salt.

6. The method of claim 5 wherein said cGMP PDE9 inhibitor is a compound of the formula (I)



20


or a pharmaceutically acceptable salt, solvate or prodrug thereof,
 wherein:

R^1 is H or (C_1-C_6) alkyl;

R^2 is (C_1-C_6) alkyl, straight chain or branched chain, (C_3-C_7) cycloalkyl or heteroaryl;

25 R^3 is (C_1-C_6) alkyl, straight chain or branched chain, optionally substituted by 1-2 groups each independently selected from Ar, (C_3-C_7) cycloalkyl, OAr, SAR, $NC(O)(C_1-C_6)$ alkyl, heteroaryl, xanthene, and naphthalene;

Ar is a group of formula



30 wherein R^4 , R^5 and R^6 are each independently selected from H, halo, phenoxy, phenyl,


- CF₃, OCF₃, S(C₁-C₈)alkyl, (C₁-C₈)alkyl, O(C₁-C₈)alkyl, said alkyl optionally substituted by a heteroaryl group or by a phenyl group, wherein said phenyl group is optionally substituted by 1-3 groups selected from halo, CF₃, OCF₃ and (C₁-C₈)alkyl; or wherein R⁴ and R⁵ may combine to form a (C₂-C₃)alkyl link, wherein said link may optionally incorporate a
- 5 heteroatom selected from O, S and N; and
- heteroaryl is aromatic 5-6 membered heterocycle containing 1-3 heteroatoms, each independently selected from O, S and N, said heterocycle optionally substituted by 1-3 substituents, each independently selected from (C₁-C₈)alkyl, halo and phenyl, said phenyl optionally substituted by 1-3 groups selected from halo and (C₁-C₈)alkyl;
- 10 with the proviso that when R¹ is -CH₃, R² cannot be -CH₂CH₂CH₃.

7. The method of claim 6 wherein R¹ is H or CH₃; R² is (C₃-C₄)alkyl, cyclopentyl or pyridinyl; R³ is (C₁-C₃)alkyl, optionally substituted by 1-2 groups selected from Ar, (C₃-C₇)cycloalkyl and heteroaryl; R⁴, R⁵ and R⁶ are each independently selected from H, halo, phenoxy, phenyl, CF₃, OCF₃, S(C₁-C₈)alkyl, (C₁-C₈)alkyl, O(C₁-C₈)alkyl; said
- 15 alkyl in the definition of R⁴, R⁵ and R⁶ is optionally substituted by a heteroaryl group or by a phenyl group optionally substituted by 1-3 groups selected from halo, CF₃, OCF₃ and (C₁-C₈)alkyl; or wherein R⁴ and R⁵ may combine to form a C₂ alkyl link, said link incorporating an O atom; and heteroaryl is an aromatic 5-6 membered heterocycle containing at least 2 nitrogen atoms, said heterocycle optionally substituted by 1-3
- 20 substituents, each independently selected from (C₁-C₈)alkyl, halo and phenyl, said phenyl in the definition of heterocycle optionally substituted by 1-3 groups selected from halo and (C₁-C₈)alkyl.

8. The method of claim 5 comprising administering to said mammal 5-(3-chloro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one, a prodrug thereof
- 25 or a pharmaceutically acceptable salt thereof or of said prodrug.

9. A method of treating dyslipidemia in a mammal comprising administering to said mammal a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said PDE9 inhibitor, prodrug, solvate or salt.

10. The method of claim 9 wherein said cGMP PDE9 inhibitor is a compound
- 30 of the formula (I)



or a pharmaceutically acceptable salt, solvate or prodrug thereof,

wherein:

R^1 is H or (C_1-C_6) alkyl;

R^2 is (C_1-C_6) alkyl, straight chain or branched chain, (C_3-C_7) cycloalkyl or heteroaryl;

R^3 is (C_1-C_6) alkyl, straight chain or branched chain, optionally substituted by 1-2 groups

- 5 each independently selected from Ar, (C_3-C_7) cycloalkyl, OAr, SAr, $NC(O)(C_1-C_6)$ alkyl, heteroaryl, xanthene, and naphthalene;


Ar is a group of formula



- wherein R^4 , R^5 and R^6 are each independently selected from H, halo, phenoxy, phenyl, CF_3 , OCF_3 , $S(C_1-C_6)$ alkyl, (C_1-C_6) alkyl, $O(C_1-C_6)$ alkyl, said alkyl optionally substituted by a heteroaryl group or by a phenyl group, wherein said phenyl group is optionally substituted by 1-3 groups selected from halo, CF_3 , OCF_3 and (C_1-C_6) alkyl; or wherein R^4 and R^5 may combine to form a (C_2-C_3) alkyl link, wherein said link may optionally incorporate a heteroatom selected from O, S and N; and
- 15 heteroaryl is aromatic 5-6 membered heterocycle containing 1-3 heteroatoms, each independently selected from O, S and N, said heterocycle optionally substituted by 1-3 substituents, each independently selected from (C_1-C_6) alkyl, halo and phenyl, said phenyl optionally substituted by 1-3 groups selected from halo and (C_1-C_6) alkyl; with the proviso that when R^1 is $-CH_3$, R^2 cannot be $-CH_2CH_2CH_3$.

- 20 11. A combination comprising a cGMP PDE9 inhibitor and one or more of a protein kinase inhibitor; an AMP-activated protein kinase; a weight loss agent; insulin; a PPAR- γ agonist; a PPAR- γ antagonist; a PPAR- α agonist; a dual PPAR- γ /PPAR- α agonist; a sorbitol dehydrogenase inhibitor; a glycogen phosphorylase inhibitor; a biguanide; an HMG-CoA reductase inhibitor; an aldose reductase inhibitor; or a PDE5
- 25 inhibitor.

12. The combination of claim 11 wherein said cGMP PDE9 inhibitor is a compound of the formula (I)



or a pharmaceutically acceptable salt, solvate or prodrug thereof,

- 30 wherein:


R¹ is H or (C₁-C₆)alkyl;

R² is (C₁-C₆)alkyl, straight chain or branched chain, (C₃-C₇)cycloalkyl or heteroaryl;

R³ is (C₁-C₆)alkyl, straight chain or branched chain, optionally substituted by 1-2 groups each independently selected from Ar, (C₃-C₇)cycloalkyl, OAr, SAr, NC(O)(C₁-C₆)alkyl,

5 heteroaryl, xanthene, and naphthalene;

Ar is a group of formula



wherein R⁴, R⁵ and R⁶ are each independently selected from H, halo, phenoxy, phenyl, CF₃, OCF₃, S(C₁-C₆)alkyl, (C₁-C₆)alkyl, O(C₁-C₆)alkyl, said alkyl optionally substituted by a
10 heteroaryl group or by a phenyl group, wherein said phenyl group is optionally substituted by 1-3 groups selected from halo, CF₃, OCF₃ and (C₁-C₆)alkyl; or wherein R⁴ and R⁵ may combine to form a (C₂-C₃)alkyl link, wherein said link may optionally incorporate a heteroatom selected from O, S and N; and

heteroaryl is aromatic 5-6 membered heterocycle containing 1-3 heteroatoms, each
15 independently selected from O, S and N, said heterocycle optionally substituted by 1-3 substituents, each independently selected from (C₁-C₆)alkyl, halo and phenyl, said phenyl optionally substituted by 1-3 groups selected from halo and (C₁-C₆)alkyl; with the proviso that when R¹ is -CH₃, R² cannot be -CH₂CH₂CH₃.

13. The combination of claim 12 wherein R¹ is H or CH₃; R² is (C₃-C₄)alkyl, cyclopentyl or pyridinyl; R³ is (C₁-C₃)alkyl, optionally substituted by 1-2 groups selected
20 from Ar, (C₃-C₇)cycloalkyl and heteroaryl; R⁴, R⁵ and R⁶ are each independently selected from H, halo, phenoxy, phenyl, CF₃, OCF₃, S(C₁-C₆)alkyl, (C₁-C₆)alkyl, O(C₁-C₆)alkyl; said alkyl in the definition of R⁴, R⁵ and R⁶ is optionally substituted by a heteroaryl group or by a phenyl group optionally substituted by 1-3 groups selected from halo, CF₃, OCF₃ and
25 (C₁-C₆)alkyl; or wherein R⁴ and R⁵ may combine to form a C₂ alkyl link, said link incorporating an O atom; and heteroaryl is an aromatic 5-6 membered heterocycle containing at least 2 nitrogen atoms, said heterocycle optionally substituted by 1-3 substituents, each independently selected from (C₁-C₆)alkyl, halo and phenyl, said phenyl in the definition of heterocycle optionally substituted by 1-3 groups selected from halo and
30 (C₁-C₆)alkyl.

14. The combination of claim 13 wherein said cGMP PDE9 inhibitor is 5-(3-chloro-benzyl)-3-isopropyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one, a prodrug thereof or a pharmaceutically acceptable salt thereof or of said prodrug, and a pharmaceutically acceptable vehicle, carrier or diluent.

15. A kit comprising:
- a) a first unit dosage form comprising a cGMP PDE9 inhibitor, a prodrug or solvate thereof or a pharmaceutically acceptable salt of said compound, prodrug or solvate and a pharmaceutically acceptable carrier, vehicle or diluent;
 - 5 b) a second unit dosage form comprising:
 - a protein kinase inhibitor;
 - an AMP-activated protein kinase;
 - a weight loss agent;
 - insulin;
 - 10 a PPAR- γ agonist;
 - a PPAR- γ antagonist;
 - a PPAR- α agonist;
 - a dual PPAR- γ /PPAR- α agonist;
 - a sorbitol dehydrogenase inhibitor;
 - 15 a glycogen phosphorylase inhibitor;
 - a biguanide;
 - an HMG-CoA reductase inhibitor;
 - an aldose reductase inhibitor; or
 - a PDE5 inhibitor;
 - 20 a prodrug or solvate of said protein kinase inhibitor, AMP-activated protein, weight loss agent, insulin, PPAR- γ agonist, PPAR- γ antagonist, PPAR- α agonist, dual PPAR- γ /PPAR- α agonist, sorbitol dehydrogenase inhibitor, glycogen phosphorylase inhibitor, biguanide, vastatin, aldose reductase inhibitor or PDE5 inhibitor; or a pharmaceutically acceptable salt thereof or of said prodrug or solvate and a pharmaceutically acceptable carrier,
 - 25 vehicle or diluent; and
 - c) a container.

INTERNATIONAL SEARCH REPORT

Inter al Application No
PCT/IB 02/03754

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61P3/06 A61P3/08 A61P3/10 A61P9/10 A61P9/12
A61K31/519 A61K45/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, BIOSIS, MEDLINE, EMBASE, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6 255 456 B1 (GOODING DOUGLAS H ET AL) 3 July 2001 (2001-07-03) column 19, line 58-65 ---	5
A	EP 0 911 333 A (PFIZER LTD ;PFIZER (US)) 28 April 1999 (1999-04-28) cited in the application the whole document ---	1-15
A	EP 0 771 799 A (BAYER AG) 7 May 1997 (1997-05-07) the whole document ---	1-15
A	US 4 666 908 A (HAMILTON HARRIET W) 19 May 1987 (1987-05-19) column 2, line 30 column 10, line 43-45 claims ---	1-15
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

4 December 2002

Date of mailing of the international search report

19/12/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Pacreu Largo, M

INTERNATIONAL SEARCH REPORT

Inter al Application No
PCT/IB 02/03754

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SODERLING S H ET AL: "Identification and characterization of a novel family of cyclic nucleotide phosphodiesterase" JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, US, vol. 273, no. 25, 19 June 1998 (1998-06-19), pages 15553-15558, XP002127167 ISSN: 0021-9258 abstract page 15557 ----	1-15
A	DOUSA T P: "Cyclic-3',5'-nucleotide phosphodiesterase isozymes in cell biology and pathophysiology of the kidney" KIDNEY INTERNATIONAL, NEW YORK, NY, US, vol. 55, no. 1, 1999, pages 29-62, XP002169142 ISSN: 0085-2538 page 39 ----	1-15
P,A	WO 02 13798 A (KOPPIKER NANDAN PARMANAND ;PFIZER LTD (GB); PFIZER (US); FRYBURG D) 21 February 2002 (2002-02-21) the whole document -----	1-15

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB 02/03754

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: —
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.: —
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 1 to 10 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.1

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

Continuation of Box I.2

Present claims 1, 5, 9, 11 and 15 relate to a compound defined by reference to a desirable characteristic or property, namely "cGMP PDE9 inhibitor". This also applies to the second active agents as defined in claims 11 and 15 which may be combined with the "cGMP PDE9 inhibitor".

The claims cover all compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds.

Furthermore, the expression "prodrug" in claims 1, 2, 4, 5, 6, 8, 9, 10, 12, 14 and 15 relates to an extremely large number of possible compounds. Support and/or disclosure within the meaning of Articles 5 and 6 PCT could not be found.

In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compound by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the compounds as defined by the general formula (I) in claims 2, 6, 10 and 12 and the specific compound of claims 4, 8 and 14.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Inter national Application No

PCT/IB 02/03754

Patent cited in search report	Publication date	Patent family member(s)	Publication date
US 625 456 B1	03-07-2001	US 5922595 A	13-07-1999
		US 2002007046 A1	17-01-2002
		AU 1710899 A	28-06-1999
		CA 2313451 A1	17-06-1999
		EP 1038003 A1	27-09-2000
		JP 2001526036 T	18-12-2001
		WO 9929873 A1	17-06-1999
EP 0911333 A	28-04-1999	AT 215954 T	15-04-2002
		BR 9804214 A	14-12-1999
		CA 2251453 A1	24-04-1999
		DE 69804743 D1	16-05-2002
		DE 69804743 T2	08-08-2002
		DK 911333 T3	01-07-2002
		EP 0911333 A1	28-04-1999
		JP 3270830 B2	02-04-2002
		JP 11217383 A	10-08-1999
		PT 911333 T	30-08-2002
		US 6235742 B1	22-05-2001
EP 0771799 A	07-05-1997	DE 19541264 A1	07-05-1997
		AU 720305 B2	25-05-2000
		AU 7053296 A	15-05-1997
		BR 9605434 A	04-08-1998
		CA 2189401 A1	07-05-1997
		CZ 9603250 A3	11-06-1997
		EE 9600172 A	16-06-1997
		EP 0771799 A1	07-05-1997
		HU 9603071 A2	28-05-1997
		JP 9124648 A	13-05-1997
		NO 964667 A	07-05-1997
		NZ 299683 A	24-09-1998
		PL 316816 A1	12-05-1997
		SG 64953 A1	25-05-1999
		SK 142596 A3	07-05-1997
		TR 970424 A2	21-05-1997
		US 5861396 A	19-01-1999
		ZA 9609278 A	02-06-1997
US 4666908 A	19-05-1987	EP 0201188 A2	17-12-1986
		JP 61236778 A	22-10-1986
WO 0213798 A	21-02-2002	AU 7660701 A	25-02-2002
		WO 0213798 A2	21-02-2002
		US 2002165237 A1	07-11-2002
		WO 02060422 A2	08-08-2002
		US 2002143015 A1	03-10-2002

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)